



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Ethambutol-d₄ dihydrochloride

Cat. No.:	HY-B0535AS
Molecular Formula:	C ₁₀ H ₂₂ D ₄ Cl ₂ N ₂ O ₂
Molecular Weight:	281.26
Target:	Antibiotic; Bacterial; Isotope-Labeled Compounds
Pathway:	Anti-infection; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Ethambutol-d ₄ dihydrochloride is deuterated labeled Ethambutol dihydrochloride (HY-B0535A). Ethambutol dihydrochloride is an antimycobacterial compound that prevents cell wall formation by inhibiting arabinosyltransferase activity.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Ethambutol dihydrochloride can be used in animal modeling to build a model of hyperuricemia. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Ethambutol. Tuberculosis (Edinb), 2008. 88(2): p. 102-5.
- [2]. Rastogi, N., V. Labrousse, and K.S. Goh, In vitro activities of fourteen antimicrobial agents against drug susceptible and resistant clinical isolates of Mycobacterium tuberculosis and comparative intracellular activities against the virulent H37Rv strain in human macrophages. Curr Microbiol, 1996. 33(3): p. 167-75.
- [3]. Kaur, D. and G.K. Khuller, In vitro, ex-vivo and in vivo activities of ethambutol and sparfloxacin alone and in combination against mycobacteria. Int J Antimicrob Agents, 2001. 17(1): p. 51-5.
- [4]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA