



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Bicyclol

Cat. No.:	HY-B0766
CAS No.:	118159-48-1
Molecular Formula:	C ₁₉ H ₁₈ O ₉
Molecular Weight:	390.34
Target:	HBV; Autophagy
Pathway:	Anti-infection; Autophagy
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (256.19 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	2.5619 mL	12.8093 mL	25.6187 mL
		5 mM	0.5124 mL	2.5619 mL	5.1237 mL
		10 mM	0.2562 mL	1.2809 mL	2.5619 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.40 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2.5 mg/mL (6.40 mM); Suspended solution; Need ultrasonic				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.40 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Bicyclol(SY 801) is a anti-hepatitis drug.Target: HBVOral administration of bicyclol normalized the elevated serum transaminases (ALT, AST) by approximately 50% in chronic viral hepatitis B and C, and also showed certain level of inhibiting HBV and HCV replication. No noticeable adverse reaction has been observed. In combination therapy of bicyclol with interferon alpha, lamivudine and adefovir dipivoxil in HBV or HCV, bicyclol may potentiate the anti-viral efficacy and reduce YMDD mutant and side effects. In 2004 China FDA issued license to manufacture bicyclol. Since then bicyclol has been widely used to treat chronic HBV and HCV in China [1].
-------------	--

REFERENCES

[1]. Walsh, S.W., Y. Wang, and A. Killian, AA-2414, an antioxidant and thromboxane receptor blocker, completely inhibits peroxide-induced vasoconstriction in the human placenta. J Pharmacol Exp Ther, 1999. 290(1): p. 220-6.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA