



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Fenipentol

Cat. No.:	HY-B1273
CAS No.:	583-03-9
Molecular Formula:	C ₁₁ H ₁₆ O
Molecular Weight:	164.25
Target:	Fluorescent Dye
Pathway:	Others
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (608.84 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	6.0884 mL	30.4418 mL	60.8835 mL
		5 mM	1.2177 mL	6.0884 mL	12.1767 mL
		10 mM	0.6088 mL	3.0442 mL	6.0884 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (15.22 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (15.22 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil				
	Solubility: ≥ 2.5 mg/mL (15.22 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Fenipentol (1-Phenyl-1-pentanol), a synthetic derivative of an ingredient of Curcuma longa that is used as a condiment and dye. Fenipentol is also an orally active choleric agent that plays an important role in release of secretin, gastrin, and pancreatic secretion of bicarbonate and protein ^{[1][2][3]} .
In Vivo	Fenipentol (PHP) (50-200 mg/kg; p.o. and intraduodenal administration) increases the secretory volume of pancreatic juice and the output of protein in rats^[2]. Fenipentol (50-200 mg/kg for i.p. and 5-10 mg/kg for i.v.) considerably increases biliary secretion in rats^[2]. Fenipentol (25-200 mg/kg; intraduodenal administration) stimulates the pancreatic secretion in the canine^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Fregnan GB, et, al. Therapeutic properties of dihydroxy-dibutylether on sub-acute liver damage induced by several hepatotoxic agents in rats. *Int J Tissue React.* 1982; 4(4): 309-18.
- [2]. Wakabayashi T, et, al. [Effect of 1-phenyl-1-hydroxy-n-pentane on pancreatic secretion in the rat and the dog]. *Nihon Yakurigaku Zasshi.* 1971 Nov; 67(6): 637-47.
- [3]. Sharma A, et, al. Biocatalytic and electrochemical reduction of selected phenyl alkanones with antibacterial activity of the corresponding phenyl alkanols. *Ijabpt Com,* 2011.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA