



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



N-Ethylmaleimide

Cat. No.:	HY-D0843
CAS No.:	128-53-0
Molecular Formula:	C ₆ H ₇ NO ₂
Molecular Weight:	125.13
Target:	Cathepsin; Deubiquitinase; Apoptosis
Pathway:	Metabolic Enzyme/Protease; Cell Cycle/DNA Damage; Apoptosis
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 50 mg/mL (399.58 mM; Need ultrasonic) DMSO : 50 mg/mL (399.58 mM; Need ultrasonic) Ethanol : 12.5 mg/mL (99.90 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	7.9917 mL	39.9584 mL	79.9169 mL
		5 mM	1.5983 mL	7.9917 mL	15.9834 mL
		10 mM	0.7992 mL	3.9958 mL	7.9917 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: PBS Solubility: 100 mg/mL (799.17 mM); Clear solution; Need ultrasonic				
	2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (16.62 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (16.62 mM); Clear solution				
	4. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (16.62 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	N-Ethylmaleimide (NEM) derives from maleic acid, it can alkylates free sulfhydryl. N-Ethylmaleimide is an irreversible cysteine protease inhibitor. N-ethylmaleimide specific inhibits phosphate transport in mitochondria. N-Ethylmaleimide inhibits prolyl endopeptidase with an IC ₅₀ value of 6.3 μM. N-Ethylmaleimide can be used to modify cysteine residues in proteins and peptides ^{[1][2][3]} .
-------------	---

IC ₅₀ & Target	IC50: 6.3 μM (prolyl endopeptidase) ^[2]	
In Vitro	N-Ethylmaleimide (20 μM;30 min) inhibits Akt Ser-473, Akt Thr-308 , p70S6K, ribosomal protein S6, 4E-BP1, eIF4E, BAD and FKHR-L1 phosphorylation ^[2] .	
	N-Ethylmaleimide (20 μM;30 min) affects conversion of pro-caspase-3 in vascular smooth muscle cells ^[2] .	
	N-Ethylmaleimide (20 μM;6 h) promotes vascular smooth muscle cells apoptosis ^[2] .	
	N-Ethylmaleimide (20 μM;30 min) affects PP2A activity and ROS production in vascular smooth muscle cells ^[2] .	
	MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Western Blot Analysis ^[2]	
	Cell Line:	Vascular smooth muscle cells
	Concentration:	20 μM
	Incubation Time:	2 hours
	Result:	Effectively inhibited platelet-derived growth factor-BB (PDGF-BB)-stimulated Akt Ser-473 , Akt Thr-308, p70S6K, ribosomal protein S6, 4E-BP1, BAD and FKHR-L1 phosphorylation with a concentration of 20 μM.
	Western Blot Analysis ^[2]	
	Cell Line:	Vascular smooth muscle cells
	Concentration:	20 μM
	Incubation Time:	2 hours
	Result:	Increased of 1.8-fold in the conversion of pro-caspase-3 into active form, and showed better effect with 20 ng/ml PDGF-BB adding.
	Apoptosis Analysis ^[2]	
Cell Line:	Vascular smooth muscle cells	
Concentration:	20 μM	
Incubation Time:	6 hours	
Result:	Induced vascular smooth muscle cells apoptosis by 3-fold, and exhibited 5-fold apoptosis with 20 ng/ml PDGF-BB adding.	
Cell Viability Assay ^[2]		
Cell Line:	Vascular smooth muscle cells	
Concentration:	20 μM	
Incubation Time:	30 min	
Result:	Increased PP2A activity 1.7-flod and increased ROS production 2-fold in vascular smooth muscle cells.	
In Vivo	N-Ethylmaleimide (10 mg/kg; i.h.) promotes the prevalence situation of mice with acute gastric ulcers ^[3] .	
	MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Animal Model:	Male Wistar rats with acute gastric ulcers induced by absolute ethanol injection ^[3]

Dosage:	10 mg/kg
Administration:	Subcutaneous injection; 10 mg/kg once
Result:	Increased the lesion area of acute gastric ulcers and attenuated the gastroprotective effect of PAG in rats.

CUSTOMER VALIDATION

- Adv Sci (Weinh). 2023 Jan 15;e2203869.
- J Exp Clin Cancer Res. 2023 Mar 30;42(1):77.
- Clin Transl Med. 2023 Jul;13(7):e1333.
- Cell Chem Biol. 2021 Nov 23;S2451-9456(21)00482-7.
- Cell Death Discov. 2022 Mar 9;8(1):107.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Moriyama A, et al. Porcine muscle prolyl endopeptidase and its endogenous substrates. J Biochem. 1988 Jul;104(1):112-7.
- [2]. Yellaturu CR, et al. N-Ethylmaleimide inhibits platelet-derived growth factor BB-stimulated Akt phosphorylation via activation of protein phosphatase 2A. J Biol Chem. 2002 Oct 18;277(42):40148-55.
- [3]. Matsuda H, et al. Roles of capsaicin-sensitive sensory nerves, endogenous nitric oxide, sulfhydryls, and prostaglandins in gastroprotection by momordin Ic, an oleanolic acid oligoglycoside, on ethanol-induced gastric mucosal lesions in rats. Life Sci. 1999;65(2):PL27-32.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA