



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Glaucocalyxin B

Cat. No.:	HY-N2113
CAS No.:	80508-81-2
Molecular Formula:	C <sub>22</sub> H <sub>30</sub> O <sub>5</sub>
Molecular Weight:	374.47
Target:	Autophagy
Pathway:	Autophagy
Storage:	<div>Powder -20°C 3 years</div> <div>In solvent -80°C 6 months</div> <div>-20°C 1 month</div>



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (267.04 mM; Need ultrasonic)					
	Preparing Stock Solutions	<div><div>Solvent</div><div>Concentration</div></div>	Mass	1 mg	5 mg	10 mg
		1 mM		2.6704 mL	13.3522 mL	26.7044 mL
		5 mM		0.5341 mL	2.6704 mL	5.3409 mL
		10 mM		0.2670 mL	1.3352 mL	2.6704 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1.25 mg/mL (3.34 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1.25 mg/mL (3.34 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 1.25 mg/mL (3.34 mM); Clear solution					

### BIOLOGICAL ACTIVITY

Description	Glaucocalyxin B is an ent kaurane diterpenoid isolated from the Chinese traditional medicine <i>Rabdosia japonica</i> with anticancer and antitumor activity; decreases the growth of HL-60 cells with an IC <sub>50</sub> of approximately 5.86 μM at 24 h.
IC <sub>50</sub> & Target	IC <sub>50</sub> : 5.86 μM (HL-60 cell Growth) <sup>[1]</sup>
In Vitro	Glaucocalyxin A (GlnA) and (GlnB) dose-dependently decrease the growth of HL-60 cells with an IC <sub>50</sub> of approximately 6.15 and 5.86 μM at 24 h, respectively. Both Gln A and B could induce apoptosis, G2/M-phase cycle arrest, DNA damage and the accumulation of reactive oxygen species (ROS) in HL-60 cells <sup>[1]</sup> . GlnB inhibits the proliferation of human cervical cancer cells

in vitro through the induction of apoptosis and autophagy, which may be mediated by the phosphatidylinositol 4,5 bisphosphate 3 kinase/Akt signaling pathway. Treatment with GlnB inhibits the proliferation of HeLa and SiHa cervical cancer cell lines in a dose dependent manner. GlnB increases the apoptotic cell population of and enhanced poly (ADP ribose) polymerase 1 cleavage. GlnB also induces increased light chain 3 II/I protein cleavage, indicating the induction of autophagy. GlnB treatment increases the expression of phosphatase and tensin homolog and decreases the expression of phosphorylated protein kinase B<sup>[2]</sup>. Glaucocalyxin B (GLB), one of five ent-kauranoid diterpenoids, significantly decreased the generation of nitric oxide (NO), tumor necrosis factor (TNF)- $\alpha$ , interleukin (IL)-1 $\beta$ , cyclooxygenase (COX)-2 and inducible nitric oxide synthase (iNOS) in the lipopolysaccharide (LPS)-activated microglia cells<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Cell Assay <sup>[3]</sup>

The microglia cells viability is assessed by MTT assay. Cells are seeded in 96-well plates at the density of  $5 \times 10^4$  cells/well. The cell culture supernatant is discarded after treatment with various agents, and then 30  $\mu$ L of MTT (0.5 mg/mL) solution is added into each well. After incubation for 4 h at 37 °C, 100  $\mu$ L of DMSO is added into each well to dissolve the formazan dye, and then the absorbance of solubilized formazan is measured by microplate reader<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## REFERENCES

- [1]. Yang WH, et al. Glaucocalyxin A and B-induced cell death is related to GSH perturbation in human leukemia HL-60 cells. *Anticancer Agents Med Chem*. 2013 Oct;13(8):1280-90.
- [2]. Pan Y, et al. Glaucocalyxin B induces apoptosis and autophagy in human cervical cancer cells. *Mol Med Rep*. 2016 Aug;14(2):1751-5.
- [3]. Gan P, et al. Anti-inflammatory effects of glaucocalyxin B in microglia cells. *J Pharmacol Sci*. 2015 May;128(1):35-46.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA