



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Kushenol A

Cat. No.:	HY-N2278
CAS No.:	99217-63-7
Molecular Formula:	C ₂₅ H ₂₈ O ₅
Molecular Weight:	408.49
Target:	Tyrosinase; Glycosidase
Pathway:	Metabolic Enzyme/Protease
Storage:	4°C, protect from light
	* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (244.80 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.4480 mL	12.2402 mL	24.4804 mL
	5 mM		0.4896 mL	2.4480 mL	4.8961 mL
	10 mM		0.2448 mL	1.2240 mL	2.4480 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Kushenol A (Leachianone E) is isolated from the root of *Sophora flavescens*. Kushenol A is a non-competitive tyrosinase inhibitor to block the conversion of L-tyrosine to L-DOPA, shows IC₅₀ and K_i values of 1.1 μM and 0.4 μM, respectively^[1]. Kushenol A is a flavonoid antioxidant, has inhibitory effects on alpha-glucosidase (IC₅₀: 45 μM; K_i: 6.8 μM) and β-amylase^[2]. Kushenol A is confirmed as potential inhibitors of enzymes targeted by cosmetics for skin whitening and aging^[1].

IC₅₀ & Target

IC₅₀: 1.1 μM (tyrosinase); 45 μM (alpha-glucosidase)^{[1][2]}

In Vitro

Kushenol A (25 μM) exhibits a highly potent inhibitory activity on ABTS+• radical scavenging with an IC₅₀ value of 9.77±0.12 μM, and exhibits inhibits scavenging activity as a percentage 93.7% at 25 μM^[1].

Kushenol A (0-60 μg/ml; 24 hours) shows considerable cytotoxic effects against NSCLC cells, exhibits IC₅₀ values of 5.3 μg/ml and 20.5 μg/ml for A549 and NCI-H226 cells, respectively. It shows an IC₅₀ value of 57.2 μg/ml for BEAS-2B cells^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Viability Assay^[3]

Cell Line:	NSCLC cell lines, A549 and NCI-H226 Normal human lung epithelial: BEAS-2B
------------	--

Concentration:	0-60 µg/ml
Incubation Time:	24 hours
Result:	Exhibited considerable cytotoxic effects against NSCLC cells.

CUSTOMER VALIDATION

- Cancer Med. 2022 Jul 4.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Kim JH, et al. Kushenol A and 8-prenylkaempferol, tyrosinase inhibitors, derived from *Sophora flavescens*. J Enzyme Inhib Med Chem. 2018 Dec;33(1):1048-1054.
- [2]. Kim JH, et al. Glycosidase inhibitory flavonoids from *Sophora flavescens*. Biol Pharm Bull. 2006 Feb;29(2):302-5.
- [3]. Chen H, et al.
A Novel Flavonoid Kushenol Z from *Sophora flavescens* Mediates mTOR Pathway by Inhibiting Phosphodiesterase and Akt Activity to Induce Apoptosis in Non-Small-Cell Lung Cancer Cells. Molecules. 2019 Dec 4;24(24). pii: E4425.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA