



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Melanotan I

Cat. No.:	HY-N2466
CAS No.:	75921-69-6
Molecular Formula:	C ₇₈ H ₁₁₁ N ₂₁ O ₁₉
Molecular Weight:	1646.85
Sequence:	Ac-Ser-Tyr-Ser-[Nle]-Glu-His-[d-Phe]-Arg-Trp-Gly-Lys-Pro-Val-NH ₂
Sequence Shortening:	Ac-SYS-[Nle]-EH-[d-Phe]-RWGKPV-NH ₂
Target:	Melanocortin Receptor
Pathway:	GPCR/G Protein; Neuronal Signaling
Storage:	Sealed storage, away from moisture



Powder -80°C 2 years
 -20°C 1 year

* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : ≥ 50 mg/mL (30.36 mM) * "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	0.6072 mL	3.0361 mL	6.0722 mL
		5 mM	0.1214 mL	0.6072 mL	1.2144 mL
		10 mM	0.0607 mL	0.3036 mL	0.6072 mL
	Please refer to the solubility information to select the appropriate solvent.				
In Vivo	1. Add each solvent one by one: PBS Solubility: 100 mg/mL (60.72 mM); Clear solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	Melanotan I is a potent non-selective melanocortin receptor (MCR) agonist. Melanotan I is a synthetic analogue of α-melanocyte stimulating hormone (α-MSH) that stimulates melanogenesis. Melanotan I can induce skin tanning by mimicking the actions of α-MSH on the melanocortin type 1 receptors (MC1R) of melanocytes. Melanotan I can be used for the research of sun-induced skin cancer, melanoma, inflammation and male erectile dysfunction ^{[1][2][3][4]} .
IC ₅₀ & Target	MC1R
In Vitro	Melanotan I (1 μM, 72 h) does not alter or slightly inhibit the formation of tumor colonies, and inhibits melanoma cell

proliferation in vitro^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Melanotan I (2 mg/kg/day for 12 weeks, s.c.) has no acute toxic effect in rodents^[2].

Melanotan I (2/10 mg/kg, s.c.) does not increase tumor incidence of human melanoma in SCID (immunodeficient) mice, nor does it lead to malignant transformation of tumors, and does not exhibit cancer-promoting effects. ^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	rats ^[2]
Dosage:	0.6 mg/kg/day for 30 days
Administration:	s.c.
Result:	Well tolerated with no change in lethality, no effect on weight gain, no serum chemistry changes, except for a slight increase (30 %) in lactic dehydrogenase levels.

CUSTOMER VALIDATION

- Theranostics. 2020 Aug 13;10(24):11110-11126.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Mahiques-Santos L. Melanotan [Melanotan]. Actas Dermosifiliogr. 2012 May;103(4):257-9. Spanish.
- [2]. Hadley ME, et al. Melanocortin peptide therapeutics: historical milestones, clinical studies and commercialization. Peptides. 2006 Apr;27(4):921-30.
- [3]. Hadley ME, et al. Discovery and development of novel melanogenic drugs. Melanotan-I and -II. Pharm Biotechnol. 1998;11:575-95.
- [4]. Langan EA, et al. Melanotropic peptides: more than just 'Barbie drugs' and 'sun-tan jabs'? Br J Dermatol. 2010 Sep;163(3):451-5.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA