



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Isodiospyrin

Cat. No.:	HY-N3488
CAS No.:	20175-84-2
Molecular Formula:	C ₂₂ H ₁₄ O ₆
Molecular Weight:	374
Target:	Topoisomerase; Bacterial; Fungal
Pathway:	Cell Cycle/DNA Damage; Anti-infection
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



BIOLOGICAL ACTIVITY

Description	Isodiospyrin, a natural dimeric naphthoquinone, is a human DNA topoisomerase I (Topoisomerase) inhibitor. Isodiospyrin can prevent both DNA relaxation and kinase activities of human topoisomerase I. Isodiospyrin shows anticancer, antibacterial and antifungal activities ^{[1][2][3]} .
IC ₅₀ & Target	Topoisomerase I
In Vitro	Isodiospyrin (10-40 μM) does not induce human topoisomerase I (htopo I)-DNA covalent complexes. However, Isodiospyrin antagonizes Camptothecin-induced, htopo I-mediated DNA cleavage. Isodiospyrin binds htopo I but not DNA. Isodiospyrin exhibits strong inhibitory effect on the kinase activity of htopo I toward splicing factor 2/alternate splicing factor in the absence of DNA^[1]. Isodiospyrin against Gram-positive bacteria with MICs ranged from 0.78 to 50 μg/mL. While Isodiospyrin against <i>Pseudomonas aeruginosa</i> ATCC 15443 and <i>S. typhi</i> ranged from 50 to 100 μg/mL. The MIC for <i>M. chelonae</i> is between 6.25 and 25 μg/mL^[2]. Isodiospyrin (30 μM; 120-144 hours) shows 81.4 % growth inhibition of <i>P. obscurans</i>. The antifungal activity of Isodiospyrin at 30 μM against <i>P. viticola</i> is 57.7 %^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Chun-Yuan Ting, et al. Isodiospyrin as a novel human DNA topoisomerase I inhibitor. *Biochem Pharmacol.* 2003 Nov 15;66(10):1981-91.
- [2]. B A Adeniyi, et al. Antibacterial activity of diospyrin, isodiospyrin and bisisodiospyrin from the root of *Diospyros piscatoria* (Gurke) (Ebenaceae). *Phytother Res.* 2000 Mar;14(2):112-7.
- [3]. Xiaoning Wang, et al. Antifungal metabolites from the roots of *Diospyros virginiana* by overpressure layer chromatography. *Chem Biodivers.* 2011 Dec;8(12):2331-40.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA