



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Benzylacetone

Cat. No.:	HY-W015616		
CAS No.:	2550-26-7		
Molecular Formula:	C ₁₀ H ₁₂ O		
Molecular Weight:	148.21		
Target:	Tyrosinase		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (674.72 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		6.7472 mL	33.7359 mL	67.4718 mL
		5 mM		1.3494 mL	6.7472 mL	13.4944 mL
		10 mM		0.6747 mL	3.3736 mL	6.7472 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (16.87 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (16.87 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (16.87 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Benzylacetone is an aromatic compound from agarwood ^[1] . Benzylacetone exhibits potent and reversible antityrosinase (mushroom) activity with IC ₅₀ s of 2.8 mM and 0.6 mM for monophenolase and diphenolase, respectively ^[2] . Benzylacetone has appetite-enhancing and locomotor-reducing effects ^[3] .
IC ₅₀ & Target	IC ₅₀ : 2.8 mM (monophenolase, tyrosinase mushroom), 0.6 mM (diphenolase, tyrosinase mushroom) ^[2]

REFERENCES

- [1]. Takamatsu S, et al. Agarotetrol: a source compound for low molecular weight aromatic compounds from agarwood heating. J Nat Med. 2018 Mar;72(2):537-541.
- [2]. Liu X, et al. Inhibition effects of benzylideneacetone, benzylacetone, and 4-phenyl-2-butanol on the activity of mushroom tyrosinase. J Biosci Bioeng. 2015 Mar;119(3):275-9.
- [3]. Ogawa K, et al. Appetite-Enhancing Effects: The Influence of Concentrations of Benzylacetone and trans-Cinnamaldehyde and Their Inhalation Time, as Well as the Effect of Aroma, on Body Weight in Mice. Biol Pharm Bull. 2016;39(5):794-8.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA