



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Methyl cinnamate-d₇

Cat. No.:	HY-W017212S
Molecular Formula:	C ₁₀ H ₃ D ₇ O ₂
Molecular Weight:	169.23
Target:	AMPK; Tyrosinase; Bacterial; Isotope-Labeled Compounds
Pathway:	Epigenetics; PI3K/Akt/mTOR; Metabolic Enzyme/Protease; Anti-infection; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Methyl cinnamate-d ₇ is deuterated labeled Cinnamyl Alcohol (HY-Y0078). Cinnamyl Alcohol is an active component from chestnut flower, inhibits increased PPAR γ expression, with anti-obesity activity ^[1] .
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1]. In 3T3-L1 cell model, Methyl cinnamate (Methyl 3-phenylpropenoate) inhibits adipocyte differentiation by attenuating expression of the adipogenic transcription factors SREBP-1, PPARγ, and C/EBPα and the transcriptional activity of PPARγ. In addition, Methyl cinnamate (Methyl 3-phenylpropenoate) activates the CaMKK2$\rightarrow$AMPK signaling cascade involved in the regulation of adipogenesis^[2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Chen YY, et al. Methyl cinnamate inhibits adipocyte differentiation via activation of the CaMKK2-AMPK pathway in 3T3-L1 preadipocytes. J Agric Food Chem. 2012 Feb 1;60(4):955-63.
- [2]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA