



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



SN-001

Cat. No.:	HY-W140974
CAS No.:	727699-84-5
Molecular Formula:	C ₂₆ H ₂₁ FN ₂ O ₄ S
Molecular Weight:	476.52
Target:	STING
Pathway:	Immunology/Inflammation
Storage:	4°C, protect from light
	* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 250 mg/mL (524.64 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.0985 mL	10.4927 mL	20.9855 mL
		5 mM		0.4197 mL	2.0985 mL	4.1971 mL
		10 mM		0.2099 mL	1.0493 mL	2.0985 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (4.36 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	SN-001 is a STING inhibitor with an IC ₅₀ of 3.82 μM ^[1] .
IC ₅₀ & Target	IC ₅₀ : 3.82 μM (STING) ^[1]
In Vitro	SN-001 targets the cyclic dinucleotide binding pocket of human STING ^[1] .
	SN-001 (5-20 μM; 6 h) significantly impairs the induction of Ifnb mRNA, in a dose-dependent manner in L929 cells ^[1] .
	SN-001 (10 μM; 3 h) inhibits cytosolic DNA-triggered STING signaling ^[1] .
	MCE has not independently confirmed the accuracy of these methods. They are for reference only.
Western Blot Analysis ^[1]	
Cell Line:	L929 cells
Concentration:	10 μM

Incubation Time:	3 h
Result:	Decreased cytosolic DNA-induced phosphorylation of STING, TBK1, IRF3, I κ B α , and p65, as well as nuclear translocation of IRF3 and p65.

REFERENCES

[1]. Hong Z, et al. STING inhibitors target the cyclic dinucleotide binding pocket. Proc Natl Acad Sci U S A. 2021 Jun 15;118(24):e2105465118.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA