



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Paclobutrazol-d₄

Cat. No.:	HY-W749242
Molecular Formula:	C ₁₅ H ₁₆ D ₄ ClN ₃ O
Molecular Weight:	297.82
Target:	Fungal
Pathway:	Anti-infection
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Paclobutrazol-d ₄ is the deuterium labeled Paclobutrazol. Paclobutrazol is a triazole-containing plant growth retardant that is known to inhibit the biosynthesis of gibberellins ^[1] .
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1]. Potential advantages of deuterated compounds: (1) Extend the half-life in vivo. Deuterated compounds may be able to prolong the pharmacokinetic characteristics of the compound, that is, prolong the half-life in vivo. This can improve compound safety, efficacy and tolerability, and increase ease of administration. (2) Improve oral bioavailability. Deuterated compounds may reduce the degree of unwanted metabolism (first-pass metabolism) in the gut wall and liver, allowing a greater proportion of the unmetabolized drug to reach its target site of action. High bioavailability determines its activity at low doses and better tolerance. (3) Improve metabolic characteristics. Deuterated compounds may reduce the formation of toxic or reactive metabolites and improve drug metabolism. (4) Improve drug safety. Deuterated compounds may reduce or eliminate adverse side effects of pharmaceutical compounds and are safe. (5) Preserve the therapeutic properties. Deuterated compounds are expected to retain similar biochemical potency and selectivity to hydrogen analogs in previous studies. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et, al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA