



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse anti ICAM-3

Catalogue number: **MUB2039P**

Clone	JA4
Isotype	IgG1/k
Product Type	Primary Antibodies
Units	0.1mg
Host	Mouse
Species reactivity	Human
Application	ELISA Flow cytometry Immunoblotting Immunocytochemistry Immunohistochemistry (frozen) Immunoprecipitation

Distributors


For Purchasing Information,
please contact your local
distributor

[Find Distributor](#)

Background

Intercellular adhesion molecule 3 (ICAM3) also known as CD50 (Cluster of Differentiation 50), is a type I transmembrane glycoprotein, which normally contain 2-9 immunoglobulin-like C2-type domains, and binds to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. It functions not only as an adhesion molecule, but also as a potent signalling molecule. ICAM-3 has previously been implicated in apoptotic cell clearance, although its precise role in the clearance process is ill defined.

Jurkat T cells that
expressed high
or low levels



Source

JA4 is a mouse monoclonal IgG1/k antibody derived by fusion of mouse Ag8.653 cells with spleen cells from a BALB/c mouse immunized with an ICAM-3/HEK transfectant.

Product

Each vial contains 100 µl 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

Applications

JA4 is suitable for immunoblotting,

immunocytochemistry, immunohistochemistry on frozen tissues and flow cytometry. Optimal antibody dilution should be determined by titration; recommended range is 1:100 – 1:200 for flow cytometry, and for immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent, and 1:100 – 1:1000 for immunoblotting applications.

Specificity

JA4 recognizes a linear epitope in domains 1 and 2 of ICAM-3.

Storage

Store at 4°C, or in small aliquots at -20°C.

References

1. Torr EE, Gardner DH, Thomas L, Goodall DM, Bielemeier A, Willetts R, Griffiths HR, Marshall LJ, Devitt A. Apoptotic cell-derived ICAM-3 promotes both macrophage chemoattraction to and tethering of apoptotic cells. *Cell Death Differ.* 2012;19: 671–679.