

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Goat anti Human Ig lambda light chains (free and bound), conjugated with FITC

Catalogue number: SR/GAHu/BJL(SD+HD)/FITC

Polyclonal
Secondary Antibodies
1 ml
Goat
Human
ELISA
Immunocytochemistry
Immunofluorescence
Immunohistochemistry (frozen)

Background

The reactivity of the antiserum is directed to the surface and hidden determinants of Ig lambda light chain. In immunoelectrophoresis this antiserum is reacting with polyclonal and monoclonal immunoglobulins of the lambda type, Bence Jones proteins as well as free light chains of the lambda type. This antiserum does not react with any other protein of human serum or plasma. This conjugate is intended for use in direct immunofluorescence for the detection of polyclonal immunoglobulins, purified monoclonal immunoglobulins of the lambda type as well as free light chains or Bence Jones proteins of the lambda type in serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Source

A pool of purified Bence Jones lambda proteins isolated from human urine. Freund's complete adjuvant is used in the first step of the immunization procedure.

Product

Fluorochrome coupled purified goat IgG lyophilized from a solution in phosphate buffered saline (pH 7.2). No preservative added, as it may interfere with the antibody activity.

Applications

Distributors

For Purchasing Information, please contact your local distributor

Find Distributor

ELISA,Immunocytochemistry,Immunohistochemistry (frozen), (In)direct immunofluorescence.

Cross Reactivity

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins and their fragments, since Ig of different species frequently share antigenic determinants. Crossreactivity of this antiSerum has not been tested in detail.

Specificity

Specificity Reference Reagent Fluorescein isothiocyanateconjugated IgG fraction of polyclonal Goat antiSerum to free and bound Human Ig lambda light chain.

Storage

The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.

Home | Company Profile | Catalogue | Distributors | Contact

Content: Nordic-MUbio BV - Copyright © 2015