

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 001-0305

Bovine IgG Fab Peroxidase

Overview

Description:	Bovine IgG Fab Fragment Peroxidase Conjugated - 001-0305
Item No.:	001-0305
Size:	1 mg
Origin:	Bovine

Product Details

Background:	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present.
Synonyms:	Bovine IgG Fab peroxidase conjugation, Bovine IgG Fab HRP conjugation, peroxidase conjugated Bovine IgG Fab, HRP conjugated Bovine IgG Fab fragment
Species of Origin:	Bovine
Conjugate:	Peroxidase (HRP)
Format:	IgG Fab
Туре:	Native Protein

Target Details

Purity/Specificity: This product was prepared from normal serum by delipidation, salt fractionation, ion exchange

chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Bovine IgG, anti-Bovine IgG F(ab')2 and anti-Bovine Serum. No reaction was observed against anti-Bovine IgG F(c) or anti-Papain.

Application Details

Application Note: Bovine IgG F(ab) fragment peroxidase conjugation can be utilized as a control or standard

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

reagent in	Western	Blotting	and	FLISA 6	experiments.
I Cug Citt iii	VVCStCIII	DIOCUITE	ullu	LLIOA	-Apcillicits.

Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be
	listed below.

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

www.rockland.com Page 3 of 3