

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 203-4302

## Chicken IgG (H&L) Antibody Peroxidase Conjugated

#### **Overview**

| Description:         | Rabbit Anti-Chicken IgG (H&L) Antibody Peroxidase Conjugated - 203-4302 |
|----------------------|---|
| Item No.:            | 203-4302  |
| Size:                | 20 mg   |
| Applications:        | ELISA   |
| Reactivity:          | Chicken   |
| <b>Host Species:</b> | Rabbit  |

#### **Product Details**

Background: Anti-Chicken IgG Peroxidase Antibody generated in rabbit detects chicken IgY. Secreted as part

of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the

antibody molecule are present.

Synonyms: rabbit anti-Chicken IgG Antibody Peroxidase Conjugated, rabbit anti-Chicken IgG HRP

Conjugated Antibody, rabbit anti-Chicken IgY Antibody Peroxidase Conjugated, rabbit anti-

Chicken IgY HRP Conjugated Antibody

Host Species: Rabbit

Specificity: IgG (H&L)

Conjugate: Peroxidase (HRP)

**IgG** 

Clonality: Polyclonal

#### **Target Details**

Immunogen:

Format:

Reactivity: Chicken

Chicken IgG whole molecule

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

**Purity/Specificity:** 

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Chicken IgG and Chicken Serum.

## **Application Details**

| <b>Tested Applications:</b> | ELISA   |
|-----------------------------|---|
| Application Note:           | Anti-Chicken IgG Peroxidase Antibody has been tested by ELISA. This product has been assayed against 1.0 ug of Chicken IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:8,000 to 1:46,000 of the reconstitution concentration is suggested for this product. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
| Assay Dilutions:            | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.   |
| ELISA:                      | 1:10,000 - 1:50,000   |
| IHC:                        | 1:500 - 1:2,500   |
| WB:                         | 1:1,000 - 1:10,000  |

#### **Formulation**

| Physical State:        | Lyophilized  |
|------------------------|--|
| Concentration:         | 10.0 mg/mL by UV absorbance at 280 nm                                  |
| Buffer:                | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2                |
| Preservative:          | None   |
| Stabilizer:            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Reconstitution Volume: | 2.0 mL   |
| Reconstitution Buffer: | Restore with deionized water (or equivalent)                           |

## **Shipping & Handling**

Shipping Condition: Ambient

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823


Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20°

C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an

undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

#### **Images**



#### **ELISA**

ELISA Results of Rabbit Anti-Chicken IgG/IgY Antibody Peroxidase Conjugated. Each well was coated in duplicate with 1.0  $\mu$ g of Chicken IgG/IgY (p/n 003-0102). The working dilution is 17,200. The starting dilution of antibody was 5 $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate (p/n TMBE-1000).

#### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3