



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Datasheet for 209-4102

Human IgG (H&L) Antibody

Overview

Description:	Rabbit Anti-Human IgG (H&L) Antibody - 209-4102
Item No.:	209-4102
Size:	50 mg
Applications:	ELISA
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Anti-Human IgG (H&L) generated in rabbit detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains — two identical heavy chains and two identical light chains arranged in a Y-shape typical of antibody monomers. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Synonyms:	Human Antibody, Rb-a-Human IgG, Human Antibody in Rabbit, Human Secondary Antibody
Host Species:	Rabbit
Specificity:	IgG (H&L)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Human
Immunogen Type:	Native Protein

Immunogen:	Anti-Human IgG whole molecule was produced by repeated immunization with Human IgG whole molecule in rabbit.
Purity/Specificity:	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Human IgG and Human Serum.

Application Details

Suggested Applications:	ELISA (Based on references)
Application Note:	Anti-Human IgG antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody based enzymatic assays requiring lot-to-lot consistency.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:100,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000

Formulation


Physical State:	Lyophilized
Concentration:	10.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	5.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
----------------------------	---------

Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images




ELISA

ELISA results using Rabbit Anti-Human IgG. Impact of behavioral variables on *M. leprae* infection levels. Anti-natural octyl disaccharide-leprosy IDRI diagnostic (NDO-LID) antibody levels in children and adolescents were measured by ELISA, using either protein A, anti- IgM or anti-IgG to detect responses. In a, samples were stratified by recorded knowledge of eating armadillo meat as either yes (n = 14) or no (n = 64). In b, samples were stratified by recorded knowledge of BCG re-vaccination following identification of the index leprosy case as either yes (n = 54) or no (n = 16). Data are displayed as box and whisker plots, with the box representing the Q1 to Q3 interquartile range and the horizontal bar representing the median of the optical density of the samples. Individual dots indicate outliers, and p-values are indicated by the lines above each indicated group. Fig. 2. PMID: 31196008.

ELISA

ELISA results using Rabbit Anti-Human IgG. Influence of index case on *M. leprae* infection levels. Anti-natural octyl disaccharide-leprosy IDRI diagnostic (NDO-LID) antibody levels in children and adolescents were measured by ELISA, using either protein A, anti- IgM or anti-IgG to detect responses. In a, samples were stratified by reported WHO operational classification of the index case as either MB (n = 66) or PB (n = 16). In b, samples were stratified by estimated duration of exposure to the index leprosy case as either less than 10 years (n = 45) or greater than 10 years (n = 37). Data are displayed as box and whisker plots, with the box representing the Q1 to Q3 interquartile range and the horizontal bar representing the median of the optical density of the samples. Individual dots indicate outliers, and p-values are indicated by the lines above each indicated group. Fig. 3. PMID: 31196008.



References

- Serrano-Coll, H et al. Social and environmental conditions related to Mycobacterium leprae infection in children and adolescents from three leprosy endemic regions of Colombia. *Bmc Infectious Diseases* (2019)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.