

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 211-1204

# Rabbit IgG F(ab')2 Antibody Fluorescein Conjugated

#### **Overview**

| Description:  | Goat Anti-Rabbit IgG F(ab')2 Antibody Fluorescein Conjugated - 211-1204 |
|---------------|---|
| Item No.:     | 211-1204  |
| Size:         | 20 mg   |
| Reactivity:   | Rabbit  |
| Host Species: | Goat  |

#### **Product Details**

Background: Anti-Rabbit IgG F(ab')2 Antibody generated in goat recognizes the dimeric Fab portion of the

rabbit IgG molecule. Rabbit IgG F(ab')2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab')2 molecules lack the Fc portion of IgG and therefore receptors that bind Rabbit IgG F(c) will not bind rabbit IgG F(ab')2 molecules. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment

composition. This Anti-Rabbit IgG F(ab')2 Antibody is conjugated to Fluorescein.

**Synonyms:** goat anti-Rabbit IgG F(ab')2 Antibody Fluorescein Conjugated, goat anti-Rabbit IgG Fab2

fragment FITC Conjugated Antibody

Host Species: Goat

**Specificity:** IgG F(ab')2

Conjugate: Fluorescein (FITC)

Clonality: Polyclonal

Format: IgG

**F/P Ratio:** 3.6

## **Target Details**

Reactivity: Rabbit

Immunogen: Rabbit IgG F(ab')2 fragment

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Purity/Specificity: This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step

process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Goat Serum, Rabbit IgG, Rabbit

IgG F(ab')2 and Rabbit Serum. No reaction was observed against Rabbit IgG F(c).

### **Application Details**

| Application Note: | This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
|-------------------|---|
| Assay Dilutions:  | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.   |
| FC:               | 1:500 - 1:2,500   |
| FLISA:            | 1:10,000 - 1:50,000   |
| IF:               | 1:1,000 - 1:5,000   |

#### **Formulation**

| Physical State:        | Lyophilized  |
|------------------------|--|
| Concentration:         | 10.0 mg/mL by UV absorbance at 280 nm                                  |
| Buffer:                | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2                |
| Preservative:          | 0.01% (w/v) Sodium Azide   |
| Stabilizer:            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Reconstitution Volume: | 2.0 mL   |
| Reconstitution Buffer: | Restore with deionized water (or equivalent)                           |

# **Shipping & Handling**

| <b>Shipping Condition:</b> | Ambient   |
|----------------------------|---|
| Storage Condition:         | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Expiration:                | Expiration date is one (1) year from date of receipt.   |

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3