

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 216-1103 Donkey IgG Fc Antibody

Overview

| Description: | Goat Anti-Donkey IgG Fc Antibody - 216-1103 |
|---------------|---|
| Item No.: | 216-1103 |
| Size: | 50 mg |
| Reactivity: | Donkey |
| Host Species: | Goat |

Product Details

| Background: | Anti-Donkey IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of donkey IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. |
|---------------|--|
| Synonyms: | goat anti-Donkey IgG F(c) fragment Antibody, goat anti-Donkey IgG Fc fragment Antibody, goat anti Donkey IgG Fc |
| Host Species: | Goat |
| Specificity: | lgG Fc |
| Clonality: | Polyclonal |
| Format: | lgG |

Target Details

| Reactivity: | Donkey |
|---------------------|--|
| Immunogen: | Donkey IgG F(c) fragment |
| Purity/Specificity: | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Donkey IgG, Donkey IgG F(c) and Donkey Serum. No reaction was observed against Donkey IgG F(ab). |



www.rockland.com tech@rockland.com +1 484.791.3823

Application Details

| Application Note: | Anti-Donkey IgG F(c) antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |
|-------------------|--|
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:20,000 - 1:100,000 |
| IHC: | 1:1,000 - 1:5,000 |
| WB: | 1:2,000 - 1:10,000 |

Formulation

| Physical State: | Lyophilized |
|-------------------------------|--|
| Concentration: | 10.0 mg/mL by UV absorbance at 280 nm |
| Buffer: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Preservative: | 0.01% (w/v) Sodium Azide |
| Stabilizer: | None |
| Reconstitution Volume: | 5.0 mL |
| Reconstitution Buffer: | Restore with deionized water (or equivalent) |

Shipping & Handling

| Shipping Condition: | Ambient |
|---------------------|---|
| Storage Condition: | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Expiration: | Expiration date is one (1) year from date of receipt. |

Disclaimer



Order online now!

www.rockland.com tech@rockland.com +1 484.791.3823

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.