

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

# Datasheet for 601-4102 Bovine IgG (H&L) Antibody

#### **Overview**

Description:	Rabbit Anti-Bovine IgG (H&L) Antibody - 601-4102
Item No.:	601-4102
Size:	2 mg
Applications:	ELISA
Reactivity:	Bovine
<b>Host Species:</b>	Rabbit

#### **Product Details**

Background:	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Anti-Bovine IgG (H&L) antibody is ideal for investigators in Immunology, Cancer, and Microbiology research.
Synonyms:	rabbit anti-Bovine IgG Antibody, Rb-a-Bovine IgG, Bovine IgG Antibody in Rabbit, rabbit anti Bovine IgG
Host Species:	Rabbit
Specificity:	IgG (H&L)
Clonality:	Polyclonal
Format:	IgG

### **Target Details**

Reactivity:	Bovine
Immunogen:	Bovine IgG whole molecule

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Purity/Specificity: Anti-Bovine IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity

chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption (s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single

precipitin arc against anti-Rabbit Serum, Bovine IgG and Bovine Serum.

### **Application Details**

<b>Tested Applications:</b>	ELISA
Application Note:	Anti-Bovine IgG (H&L) Antibody has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody based enzymatic assays requiring lot-to-lot consistency.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:100,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000

#### **Formulation**

Stabilizer:	None
Preservative:	0.01% (w/v) Sodium Azide
Buffer:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	2.0 mg/mL by UV absorbance at 280 nm
Physical State:	Liquid (sterile filtered)

### **Shipping & Handling**


Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Expiration:	Expiration date is one (1) year from date of receipt.

#### **Images**

www.rockland.com Page 2 of 3



www.rockland.com tech@rockland.com +1 484.791.3823



#### **ELISA**

ELISA Results of Purified Rabbit Anti-Bovine IgG Antibody tested against purified Bovine IgG. Each well was coated in duplicate with 1.0  $\mu$ g of Bovine IgG (p/n 001-0102). The working dilution is 1:420,000. The starting dilution of antibody was 5 $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using HRP Conjugation Stabilizer (p/n MB-076), Goat Anti-Rabbit HRP conjugated (p/n 611-103-122) and TMB substrate (p/n TMBE-1000).

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3