



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



**Datasheet for 604-4102****Dog IgG Antibody****Overview**

|                      |   |
|----------------------|---|
| <b>Description:</b>  | Rabbit Anti-Dog IgG Antibody - 604-4102 |
| <b>Item No.:</b>     | 604-4102                                |
| <b>Size:</b>         | 10 mg                                   |
| <b>Applications:</b> | ELISA                                   |
| <b>Reactivity:</b>   | Dog                                     |
| <b>Host Species:</b> | Rabbit                                  |

**Product Details**

|                      |   |
|----------------------|---|
| <b>Background:</b>   | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. Anti-Dog IgG (H&L) antibody is ideal for investigators in Immunology, Cancer, and Microbiology research. |
| <b>Synonyms:</b>     | rabbit anti-Dog IgG antibody, rabbit anti Dog IgG antibody  |
| <b>Host Species:</b> | Rabbit  |
| <b>Specificity:</b>  | IgG (H&L)   |
| <b>Clonality:</b>    | Polyclonal  |
| <b>Format:</b>       | IgG   |

**Target Details**

|                    |                        |
|--------------------|------------------------|
| <b>Reactivity:</b> | Dog                    |
| <b>Immunogen:</b>  | Dog IgG whole molecule |

**Purity/Specificity:** Anti-Dog IgG (H&L) antibody was purified from concentrated monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Dog IgG and Dog Serum.

## Application Details

|                             |   |
|-----------------------------|---|
| <b>Tested Applications:</b> | ELISA   |
| <b>Application Note:</b>    | Anti-Dog IgG (H&L) antibody has been tested by ELISA. Specific conditions for reactivity should be optimized by the end user. |
| <b>Assay Dilutions:</b>     | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.                               |
| <b>ELISA:</b>               | 1:200,000   |

## Formulation

|                        |  |
|------------------------|--|
| <b>Physical State:</b> | Liquid (sterile filtered)                                  |
| <b>Concentration:</b>  | 7.56 mg/mL by UV absorbance at 280 nm                      |
| <b>Buffer:</b>         | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| <b>Preservative:</b>   | 0.01% (w/v) Sodium Azide                                   |
| <b>Stabilizer:</b>     | None   |

## Shipping & Handling

|                            |  |
|----------------------------|--|
| <b>Shipping Condition:</b> | Wet Ice  |
| <b>Storage Condition:</b>  | Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. |
| <b>Expiration:</b>         | Expiration date is one (1) year from date of receipt.  |

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.