

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



#### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 605-4003

# **Goat IgG Fc Antibody Rhodamine Conjugated**

#### **Overview**

Description:	Rabbit Anti-Goat IgG Fc Antibody Rhodamine Conjugated - 605-4003
Item No.:	605-4003
Size:	2 mg
Reactivity:	Goat
Host Species:	Rabbit

### **Product Details**

Background:	Anti-Goat IgG F(c) generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of goat IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity.
Synonyms:	Rabbit anti-Goat IgG F(c) Antibody Rhodamine Conjugation, Rabbit anti-Goat IgG Fc fragment Rhodamine Conjugated Antibody
<b>Host Species:</b>	Rabbit
Specificity:	IgG Fc
Conjugate:	Rhodamine (TRITC)
Clonality:	Polyclonal
Format:	IgG
F/P Ratio:	4.0

### **Target Details**

Reactivity:	Goat
Immunogen:	Goat IgG F(c) fragment
Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG. Goat IgG F(c), and Goat Serum. No reaction was observed against Goat IgG F(ab).

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

## **Application Details**

Application Note:	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
FC:	1:500 - 1:2,500
FLISA:	1:10,000 - 1:50,000
IF:	1:1,000 - 1:5,000

#### **Formulation**

Concentration:       2.0 mg/mL by UV absorbance at 280 nm         Buffer:       0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2         Preservative:       0.01% (w/v) Sodium Azide         Stabilizer:       10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free         Reconstitution Volume:       1.0 mL         Reconstitution Buffer:       Restore with deionized water (or equivalent)	Physical State:	Lyophilized
Preservative:       0.01% (w/v) Sodium Azide         Stabilizer:       10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free         Reconstitution Volume:       1.0 mL	Concentration:	2.0 mg/mL by UV absorbance at 280 nm
Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free  Reconstitution Volume: 1.0 mL	Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume: 1.0 mL	Preservative:	0.01% (w/v) Sodium Azide
	Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Buffer: Restore with deionized water (or equivalent)	Reconstitution Volume:	1.0 mL
	Reconstitution Buffer:	Restore with deionized water (or equivalent)

## **Shipping & Handling**

<b>Shipping Condition:</b>	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

#### Disclaimer

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3