



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 605-4302**Goat IgG (H&L) Secondary Antibody Peroxidase Conjugated****Overview**

Description:	Rabbit Anti-Goat IgG (H&L) Antibody Peroxidase Conjugated - 605-4302
Item No.:	605-4302
Size:	2 mg
Applications:	ELISA, IHC, WB
Reactivity:	Goat
Host Species:	Rabbit

Product Details

Background:	Anti-Goat IgG Peroxidase Antibody generated in rabbit detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Synonyms:	rabbit anti-Goat IgG Peroxidase Conjugated Antibody, rabbit anti-Goat IgG HRP Conjugated Antibody, HRP anti-goat IgG, Peroxidase anti-Goat IgG
Host Species:	Rabbit
Specificity:	IgG (H&L)
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Goat
--------------------	------

Immunogen:	Goat IgG whole molecule
Purity/Specificity:	Rabbit Anti-Goat IgG secondary Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Goat IgG and Goat Serum.
Relevant Links:	<ul style="list-style-type: none">• 605-4302 SDS

Application Details

Tested Applications:	ELISA
Suggested Applications:	IHC, WB (Based on references)
Application Note:	Anti-Goat IgG Peroxidase conjugate has been tested by ELISA and western blot. This antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:50,000 - 1:100,000
IHC:	1:1,000 - 1:5,000
WB:	1:5,000 - 1:20,000

Formulation


Physical State:	Lyophilized
Concentration:	2.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
----------------------------	---------


Storage Condition:	Store Secondary Antibody at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



ELISA

ELISA results of Rabbit Anti-Goat IgG Antibody Peroxidase Conjugated tested against purified Goat IgG protein. Each well was coated in duplicate with 1.0 µg of Goat IgG (p/n 005-0102). The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. The titer is 1:22,700. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate p/n TMBE-1000.



Western Blot

Western Blot of Rabbit anti-GOAT IgG (H&L) Antibody Peroxidase Conjugated. Lane 1: Goat IgG. Load: 100 ng per lane. Secondary antibody: GOAT IgG (H&L) Antibody Peroxidase Conjugated at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT.

References

- Birnbaum SK et al. The proprotein convertase BLI-4 promotes collagen secretion prior to assembly of the *Caenorhabditis elegans* cuticle. *PLoS Genet.* (2023)

- Lim JH et al. Palmitoyl-RGD promotes the expression of dermal-epidermal junction components in HaCaT cells. *Mol Med Rep.* (2022)
- Cheng A et al. Fatty acid-binding protein 7 triggers α -synuclein oligomerization in glial cells and oligodendrocytes associated with oxidative stress. *Acta Pharmacol Sin.* (2022)
- Cheng A et al. A novel fatty acid-binding protein 5 and 7 inhibitor ameliorates oligodendrocyte injury in multiple sclerosis mouse models. *EBioMedicine* (2022)
- Cohen JD et al. A C. elegans Zona Pellucida domain protein functions via its ZPc domain. *PLoS Genet.* (2020)
- Cheng A et al. Fatty Acid Binding Protein 5 Mediates Cell Death by Psychosine Exposure through Mitochondrial Macropores Formation in Oligodendrocytes. *Biomedicines.* (2020)
- Brunet A et al. Nuclear lamin B1 interactions with chromatin during the circadian cycle are uncoupled from periodic gene expression. *Front Genet.* (2019)
- Doi T et al. Urinary IgG4 and Smad1 are specific biomarkers for renal structural and functional changes in early stages of diabetic nephropathy. *Diabetes.* (2018)
- Crater AK et al. Toxoplasma ubiquitin-like protease 1, a key enzyme in sumoylation and desumoylation pathways, is under the control of non-coding RNAs. *Int J Parasitol.* (2018)
- Hill B. A coiled-coil strategy for the directional display of multiple proteins on the surface of iron oxide nanoparticles. *Royal Society of Chemistry.* (2017)
- Haj-Yasein NN et al. Cysteine deprivation prevents induction of peroxisome proliferator-activated receptor gamma-2 and adipose differentiation of 3T3-L1 cells. *Biochim Biophys Acta Mol Cell Biol Lipids.* (2017)
- Fan et al. LXR α Regulates Hepatic ChREBP α Activity and Lipogenesis upon Glucose, but Not Fructose Feeding in Mice. *Nutrients* (2017)
- Suzuki T et al. Efficient antibody production in the methylotrophic yeast Ogataea minuta by overexpression of chaperones. *J Biosci Bioeng.* (2017)
- Gill et al. Integrity of Narrow Epithelial Tubes in the C. elegans Excretory System Requires a Transient Luminal Matrix. *PLOS Genetics* (2016)
- Storti P et al. Galectin-1 suppression delineates a new strategy to inhibit myeloma-induced angiogenesis and tumoral growth in vivo. *Leukemia.* (2016)
- Xu P et al. Gene expression levels of Casein kinase 1 (CK1) isoforms are correlated to adiponectin levels in adipose tissue of morbid obese patients and site-specific phosphorylation mediated by CK1 influences multimerization of adiponectin. *Mol Cell Endocrinol.* (2015)
- Bindesbøll et al. Liver X receptor regulates hepatic nuclear O-GlcNAc signaling and carbohydrate responsive element-binding protein activity. *Journal of Lipid Research* (2015)
- Klaric et al. A reduction in Npas4 expression results in delayed neural differentiation of mouse embryonic stem cells. *Stem Cell Research & Therapy* (2014)
- Cherry AA et al. Characterization of a homolog of DEAD-box RNA helicases in Toxoplasma gondii as a marker of cytoplasmic mRNP stress granules. *Gene.* (2014)

- Tarassishin L et al. Interleukin-1-induced changes in the glioblastoma secretome suggest its role in tumor progression. *Journal of Proteomics* (2014)
- Bischof J et al. CK1δ kinase activity is modulated by Chk1-mediated phosphorylation. *PLoS One* (2013)
- Nishimiya D et al. Identification of antibody-interacting proteins that contribute to the production of recombinant antibody in mammalian cells. *Appl Microbiol Biotechnol.* (2012)
- Ronsyn MW et al. Plasmid-based genetic modification of human bone marrow-derived stromal cells: analysis of cell survival and transgene expression after transplantation in rat spinal cord. *BMC Biotechnol.* (2007)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.