

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Datasheet for 605-442-013

Goat IgG (H&L) Antibody DyLight™ 549 Conjugated Pre-Adsorbed

Overview

Description:	Rabbit Anti-Goat IgG (H&L) Antibody DyLight™ 549 Conjugated (Min X Human Serum Proteins) - 605-442-013
Item No.:	605-442-013
Size:	100 μg
Reactivity:	Goat
Host Species:	Rabbit

Product Details

Background:	Anti-Goat IgG DyLight Antibody generated in rabbit detects goat IgG. Secreted as part of the
	adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum

immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F (ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of crossreactivity, and host-species source and fragment composition.

Synonyms:	rabbit anti-Goat IgG DyLight™ 549 Conjugated Antibody, rabbit anti-Goat IgG Antibody DyLight™ 549 Conjugation
Host Species:	Rabbit

Specificity:	IgG (H&L)
Conjugate:	DyLight™ 549
Clonality:	Polyclonal
Format:	IgG
F/P Ratio:	3.1

Target Details

www.rockland.com Page 1 of 4



Reactivity:	Goat	
Immunogen:	Goat IgG whole molecule	
Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against Human Serum Proteins. This antibody will react with heavy chains of Goat IgG and with light chains of most Goat immunoglobulins.	

Application Details

Application Note:	The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
FLISA:	>1:20,000
IF:	>1:5,000
WB:	>1:10,000

Formulation

Concentration: 1.0 mg/mL by UV absorbance at 280 nm Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Preservative: 0.01% (w/v) Sodium Azide Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Reconstitution Volume: 100 μL Reconstitution Buffer: Restore with deionized water (or equivalent)	Physical State:	Lyophilized		
Preservative: 0.01% (w/v) Sodium Azide Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Reconstitution Volume: 100 μL	Concentration:	1.0 mg/mL by UV absorbance at 280 nm		
Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Reconstitution Volume: 100 μL	Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2		
Reconstitution Volume: 100 μL	Preservative:	0.01% (w/v) Sodium Azide		
	Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free		
Reconstitution Buffer: Restore with deionized water (or equivalent)	Reconstitution Volume:	100 μL		
	Reconstitution Buffer:	Restore with deionized water (or equivalent)		

Shipping & Handling

Shipping Condition: Ambient

www.rockland.com Page 2 of 4




Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20°

C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an

undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

DyLight[™] dyes can be used for two-color Western Blot detection with low background and high signal. Anti-tubulin was detected using a DyLight[™] 549 conjugate. Anti-TNFa was detected using a DyLight[™] 649 conjugate. The image was captured using the Typhoon[™] 9410 Imaging System.

Diagram


Properties of DyLight™ Conjugates.

Emission	Color	DyLight™ Dye	Ex/Em (nm)	е (M-1 cm-1)	Similar Dyes
Blue		405	400/420	30,000	Alexa™ 405, Cascade Blue
Green		488	493/518	70,000	Alexa™ 488, Cy2®, FITC
Yellow		549	550/568	150,000	Alexa™ 546, Alexa 555, Cy3®,TRITC
Red		649	646/674	250,000	Alexa™ 647, Cy5®
Near Infrared		680	682/715	140,000	Alexa™ 680, Cy5.5®, IRDye™ 700
Infrared		800	770/794	270,000	IRDye™ 800

www.rockland.com Page 3 of 4







Diagram

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 4 of 4