

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





#### Datasheet for 605-746-002

# Goat IgG (H&L) Antibody DyLight™ 405 Conjugated

#### **Overview**

Description:	Donkey Anti-Goat IgG (H&L) Antibody DyLight™ 405 Conjugated - 605-746-002			
Item No.:	605-746-002			
Size:	100 μg			
Applications:	WB			
Reactivity:	Goat			
<b>Host Species:</b>	Donkey			

#### **Product Details**

**Background:** Anti-Goat IgG DyLight Antibody generated in donkey detects goat IgG. Secreted as part of the

adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F (ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody

molecule are present.

Synonyms: donkey anti-Goat IgG Antibody DyLight™ 405 Conjugation, donkey anti-Goat IgG DyLight™ 405

Conjugated Antibody

Host Species: Donkey

Specificity: IgG (H&L)

Conjugate: DyLight™ 405

Clonality: Polyclonal

Format: IgG

**F/P Ratio:** 2.0

### **Target Details**

Reactivity: Goat

www.rockland.com Page 1 of 4





Immunogen:	Goat IgG, whole molecule
Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by conjugation to fluorochrome and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Goat IgG and Goat Serum. This antibody will react with heavy chains of Goat IgG and with light chains of most Goat immunoglobulins.

# **Application Details**

Suggested Applications:	WB (Based on references)  The emission spectra for this DyLight™ conjugate match the principle output wavelengths of most common fluorescence instrumentation. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.		
Application Note:			
Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.			
FLISA:	>1:20,000		
IF:	>1:5,000		
WB:	>1:10,000		

### **Formulation**

Physical State:	Lyophilized			
Concentration:	1.0 mg/mL by UV absorbance at 280 nm			
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2			
Preservative:	0.01% (w/v) Sodium Azide			
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free			
Reconstitution Volume:	100 μL			
Reconstitution Buffer:	stitution Buffer: Restore with deionized water (or equivalent)			

## **Shipping & Handling**

Shipping Condition: Ambient

www.rockland.com Page 2 of 4





**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20°

C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an

undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

#### **Images**

# B 1.25 μg/ml DOX 80 μM Tan IIA 0 8 16 24 (h) P73α PUMA Noxa β-actin

#### **Western Blot**

Tan IIA exposure does not activate P73 $\alpha$ -induced inhibition of viability. (B) P73 $\alpha$ , PUMA and Noxa protein expression levels in H1299 cells following exposure to to 1.25  $\mu$ g/ml DOX or 80  $\mu$ M Tan IIA for the indicated times. Figure 4. PMID: 29207086.

Emission	Color	DyLight™ Dye	Ex/Em (nm)	е (M <sup>-1</sup> cm <sup>-1</sup> )	Similar Dyes
Blue		405	400/420	30,000	Alexa™ 405, Cascade Blue
Green	- 1	488	493/518	70,000	Alexa™ 488, Cy2®, FITC
Yellow		549	550/568	150,000	Alexa™ 546, Alexa 555, Cy3®,TRITC
Red		649	646/674	250,000	Alexa™ 647, Cy5®
Near Infrared		680	682/715	140,000	<u>Alexa™</u> 680, Cy5.5®, IRDye™ 700
Infrared		800	770/794	270,000	IRDye™ 800

#### Diagram


Properties of DyLight™ Fluorescent Dyes.

www.rockland.com Page 3 of 4





# **Diagram**DyLight™ 405 Fluorescence Spectra



#### References

Zu et al. Tan IIA inhibits H1299 cell viability through the MDM4@IAP3 signaling pathway. Molecular Medicine Reports
(2018)

#### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 4 of 4