



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for 606-1603****Guinea Pig IgG Fc Antibody Biotin Conjugated****Overview**

|                      |   |
|----------------------|---|
| <b>Description:</b>  | Goat Anti-Guinea Pig IgG Fc Antibody Biotin Conjugated - 606-1603 |
| <b>Item No.:</b>     | 606-1603  |
| <b>Size:</b>         | 2 mg  |
| <b>Reactivity:</b>   | Guinea Pig  |
| <b>Host Species:</b> | Goat  |

**Product Details**

|                      |   |
|----------------------|---|
| <b>Background:</b>   | Anti-Guinea Pig IgG F(c) Biotin generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of Guinea Pig IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. |
| <b>Synonyms:</b>     | Goat Anti-Guinea Pig IgG F(c) Antibody Biotin Conjugated, Goat Anti Guinea Pig IgG Fc Fragment Antibody Biotin Conjugated, Goat Anti Guinea Pig IgG Fc Antibody Biotin Conjugated   |
| <b>Host Species:</b> | Goat  |
| <b>Specificity:</b>  | IgG Fc  |
| <b>Conjugate:</b>    | Biotin  |
| <b>Clonality:</b>    | Polyclonal  |
| <b>Format:</b>       | IgG   |

**Target Details**

|                            |   |
|----------------------------|---|
| <b>Reactivity:</b>         | Guinea Pig  |
| <b>Immunogen:</b>          | Guinea Pig IgG F(c) fragment  |
| <b>Purity/Specificity:</b> | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Guinea Pig IgG, Guinea Pig IgG F(c) and Guinea Pig Serum. No reaction was observed against Guinea Pig IgG F(ab). |

## Application Details

|                          |   |
|--------------------------|---|
| <b>Application Note:</b> | This product has been assayed against 1.0 µg of Guinea Pig IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:35,000 to 1:150,000 of the reconstitution concentration is suggested for this product. |
| <b>Assay Dilutions:</b>  | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.   |
| <b>ELISA:</b>            | 1:20,000 - 1:100,000  |
| <b>IHC:</b>              | 1:1,000 - 1:5,000   |
| <b>WB:</b>               | 1:2,000 - 1:10,000  |

## Formulation

|                               |  |
|-------------------------------|--|
| <b>Physical State:</b>        | Lyophilized  |
| <b>Concentration:</b>         | 2.0 mg/mL by UV absorbance at 280 nm                                   |
| <b>Buffer:</b>                | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2             |
| <b>Preservative:</b>          | 0.01% (w/v) Sodium Azide   |
| <b>Stabilizer:</b>            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| <b>Reconstitution Volume:</b> | 1.0 mL   |
| <b>Reconstitution Buffer:</b> | Restore with deionized water (or equivalent)                           |

## Shipping & Handling

|                            |   |
|----------------------------|---|
| <b>Shipping Condition:</b> | Ambient   |
| <b>Storage Condition:</b>  | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| <b>Expiration:</b>         | Expiration date is one (1) year from date of receipt.   |

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.