

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 607-1302 Golden Syrian Hamster IgG (H&L) Antibody Peroxidase Conjugated

Overview

| Description: | Goat Anti-Golden Syrian Hamster IgG (H&L) Antibody Peroxidase Conjugated - 607-1302 |
|---------------|---|
| Item No.: | 607-1302 |
| Size: | 2 mg |
| Applications: | ELISA |
| Reactivity: | Golden Syrian Hamster |
| Host Species: | Goat |

Product Details

| Background: | Anti-Golden Syrian Hamster IgG Peroxidase Antibody generated in goat detects Golden Syrian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host- species source and fragment composition. |
|---------------|--|
| Synonyms: | Golden Syrian Hamster Antibody, Gt-a-Golden Syrian Hamster IgG, Golden Syrian Hamster IgG Antibody in Goat, Golden Syrian Hamster Secondary Antibody |
| Host Species: | Goat |
| Specificity: | IgG (H&L) |
| Conjugate: | Peroxidase (HRP) |
| Clonality: | Polyclonal |
| Format: | lgG |

Target Details



www.rockland.com tech@rockland.com +1 484.791.3823

| Reactivity: | Golden Syrian Hamster |
|---------------------|---|
| Immunogen Type: | Native Protein |
| Immunogen: | Anti-Golden Syrian Hamster IgG whole molecule was produced by repeated immunization with Golden Syrian Hamster IgG whole molecule in goat. |
| Purity/Specificity: | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Golden Syrian Hamster IgG and Golden Syrian Hamster Serum. Diminished reactivity will occur against Armenian Hamster IgG. |

Application Details

| Tested Applications: | ELISA |
|-----------------------------|---|
| Application Note: | Anti-Golden Syrian Hamster IgG peroxidase conjugated antibody has been tested by ELISA and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user. |
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:7,000 - 1:35,000 |
| IHC: | 1:500 - 1:2,500 |
| WB: | 1:1,000 - 1:5,000 |

Formulation

| Physical State: | Lyophilized |
|-------------------------------|--|
| Concentration: | 2.0 mg/mL by UV absorbance at 280 nm |
| Buffer: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Preservative: | 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide! |
| Stabilizer: | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Reconstitution Volume: | 1.0 mL |
| Reconstitution Buffer: | Restore with deionized water (or equivalent) |

Shipping & Handling

Shipping Condition: Ambient


www.rockland.com



www.rockland.com tech@rockland.com +1 484.791.3823

| Storage Condition: | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
|--------------------|---|
| Expiration: | Expiration date is one (1) year from date of receipt. |

Images



ELISA

ELISA results of purified Polyclonal Goat Anti-Golden Syrian Hamster IgG Antibody Peroxidase Conjugated tested against Hamster IgG. Each well was coated in duplicate with 1.0µg of antigen. The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate (p/n TMBE-1000).

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.