

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 607-1602

## Golden Syrian Hamster IgG (H&L) Antibody Biotin Conjugated

#### **Overview**

| Description:  | Goat Anti-Golden Syrian Hamster IgG (H&L) Antibody Biotin Conjugated - 607-1602 |
|---------------|---|
| Item No.:     | 607-1602  |
| Size:         | 2 mg  |
| Reactivity:   | Golden Syrian Hamster   |
| Host Species: | Goat  |

#### **Product Details**

**Background:** Anti-Golden Syrian Hamster IgG Biotin Antibody generated in goat detects Golden Syrian

Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

**Synonyms:** goat anti-Golden Syrian Hamster IgG biotin conjugated antibody, goat anti-Hamster IgG biotin

conjugated antibody

**Host Species:** Goat

Specificity: IgG (H&L)

Clonality: Polyclonal

Format: IgG

### **Target Details**

Conjugate:

**Reactivity:** Golden Syrian Hamster

Biotin

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

| Immunogen:          | Hamster IgG whole molecule   |
|---------------------|--|
| Purity/Specificity: | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Hamster IgG and Hamster Serum. |

## **Application Details**

| Application Note: | This product has been assayed against 1.0 ug of Hamster IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 20,000 to 1:60,000 of the reconstitution concentration is suggested for this product. |
|-------------------|---|
| Assay Dilutions:  | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.   |
| ELISA:            | 1:20,000 - 1:100,000  |
| IHC:              | 1:1,000 - 1:5,000   |
| WB:               | 1:2,000 - 1:10,000  |

### **Formulation**

| Physical State:        | Lyophilized  |
|------------------------|--|
| Concentration:         | 2.0 mg/mL by UV absorbance at 280 nm                                   |
| Buffer:                | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2             |
| Preservative:          | 0.01% (w/v) Sodium Azide   |
| Stabilizer:            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Reconstitution Volume: | 1.0 mL   |
| Reconstitution Buffer: | Restore with deionized water (or equivalent)                           |
|                        |  |

## **Shipping & Handling**

| Shipping Condition: | Ambient   |
|---------------------|---|
| Storage Condition:  | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Expiration:         | Expiration date is one (1) year from date of receipt.   |

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3