

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 609-101-123 Human IgG (H&L) Antibody Pre-Adsorbed

Overview

Description:	Goat Anti-Human IgG (H&L) Antibody (Min X Bv Ch Gt GP Ham Hs Ms Rb Rt & Sh Serum Proteins) - 609-101-123
Item No.:	609-101-123
Size:	1 mg
Applications:	ELISA, Microarray
Reactivity:	Human
Host Species:	Goat

Product Details

Background:	Anti-Human IgG (H&L) generated in goat detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains — two identical heavy chains and two identical light chains arranged in a Y-shape typical of antibody monomers. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host- species source and fragment composition.
Synonyms:	Goat Anti Human IgG Antibody Pre-Adsorbed, Goat Anti-Human IgG Antibody Pre-Adsorbed
Host Species:	Goat
Specificity:	IgG (H&L)
Clonality:	Polyclonal
Format:	lgG

Target Details

Reactivity:	Human
Immunogen:	Human IgG whole molecule



www.rockland.com tech@rockland.com +1 484.791.3823

Purity/Specificity:This product was prepared from monospecific antiserum by immunoaffinity chromatography
using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any
unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc
against anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against
Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Mouse, Rabbit, Rat and Sheep Serum
Proteins.

Application Details

ELISA
Microarray (Based on references)
Anti-Human IgG antibody has been tested by ELISA and is suitable for western blot and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.
All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
1:20,000 - 1:100,000
1:1,000 - 1:2,000
1:2,000 - 1:10,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.18 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Expiration:	Expiration date is one (1) year from date of receipt.

ROCKLAND

www.rockland.com tech@rockland.com +1 484.791.3823

Images



ELISA

ELISA Results of Goat Anti-Human IgG Antibody (MX Bv Ch Gt GP Ham Hs Ms Rb Rt Sh Serum Proteins) tested against purified Human IgG MX10. Each well was coated in duplicate with 1.0 µg of Human IgG (p/n 009-0102) [Red Line], Bovine IgG (p/n 001-0102) [Blue Line], Chicken IgG (003-0102), Goat IgG (p/n 005-0102) [Purple Line], Guinea Pig IgG (p/n 006-0102), Hamster IgG (p/n 007-0102), Horse IgG (p/n 008-0102), Mouse IgG (p/n 010-0102), Rabbit IgG (p/n 011-0102), Rat IgG (p/n 012-0102), Sheep IgG (p/n 013-0102). The working dilution is 1:184,500. The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using Donkey Anti-Goat IgG HRP conjugated (p/n 605-703-125) and TMB substrate (p/n TMBE-1000).

References

- Metz I, Beißbarth T, Ellenberger D, et al. Serum peptide reactivities may distinguish neuromyelitis optica subgroups and multiple sclerosis. *Neurol Neuroimmunol Neuroinflamm.* (2016)
- Bucukovski, J et al. A Multiplex Label-Free Approach to Avian Influenza Surveillance and Serology. PloS One (2015)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.