



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 609-4103

Human IgG Fc Antibody

Overview

Description:	Rabbit Anti-Human IgG Fc Antibody - 609-4103
Item No.:	609-4103
Size:	5 mg
Applications:	ELISA, WB, FC, IP
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Anti-Human IgG F(c) generated in rabbit detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of human IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(c) Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.
Synonyms:	Human IgG F(c) Antibody, Human IgG Fc Antibody, Rb-a-Human IgG Fc, Human IgG F(c) Antibody in Rabbit, rabbit anti-human IgG F(c) Secondary Antibody
Host Species:	Rabbit
Specificity:	IgG Fc
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Human
Immunogen Type:	Native Protein

Immunogen:	Anti-Human IgG F(c) fragment was produced by repeated immunization with Human IgG F(c) fragment in rabbit.
Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab).

Application Details

Tested Applications:	ELISA, WB
Suggested Applications:	FC, IP (Based on references)
Application Note:	Anti-Human IgG F(c) fragment has been tested by western blot and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:26,000-1:46,000
IHC:	1:1,000 - 1:6,000
WB:	1:3,000 - 1:30,000

Formulation


Physical State:	Liquid (sterile filtered)
Concentration:	5.014 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit anti-Human Fc antibody. Lane 1: Human Fc (p/n 009-0103). Lane 2: none. Load: 100 ng per lane. Primary antibody: Human Fc antibody at 1:1,000 for overnight at 4°C. Secondary antibody: DyLight™ 649 rabbit secondary antibody at 1:20,000 for 30 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 28 kDa, 28 kDa for Human Fc. Other band(s): Human Fc splice variants and isoforms.

References

- Ito N et al. Pivotal role of myeloid-derived suppressor cells in infection-related tumor growth. *Cancer Med.* (2024)
- Ying B et al. High-Level Production of scFv-Fc Antibody Using an Artificial Promoter System with Transcriptional Positive Feedback Loop of Transactivator in CHO Cells. *Cells.* (2023)
- Izumi H et al. Developmental synapse pathology triggered by maternal exposure to the herbicide glufosinate ammonium. *Front Mol Neurosci.* (2023)
- Kamioka Y et al. Distinct bidirectional regulation of LFA1 and $\alpha 4\beta 7$ by Rap1 and integrin adaptors in T cells under shear flow. *Cell Rep.* (2023)
- Yoshida T et al. Canonical versus non-canonical transsynaptic signaling of neuroligin 3 tunes development of sociality in mice. *Nat Commun.* (2021)
- Zheng F et al. LINE-1 vectors mediate recombinant antibody gene transfer by retrotransposition in Chinese hamster ovary cells. *Biotechnol J.* (2021)
- Yukinari Kato et al. Molecular analysis of the pathophysiological binding of the platelet aggregation-inducing factor podoplanin to the C-type lectin-like receptor CLEC-2. *Cancer Sci.* (2008)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.