

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

### Datasheet for 610-103-043

# Mouse IgG3 Antibody Peroxidase Conjugated Pre-adsorbed

#### **Overview**

| Description:         | Goat Anti-Mouse IgG3 (Gamma 3 chain) Antibody Peroxidase Conjugated (Min X Bv, Hu, and Rb Serum Proteins) - 610-103-043 |
|----------------------|---|
| Item No.:            | 610-103-043   |
| Size:                | 1 mg  |
| Applications:        | ELISA, WB   |
| Reactivity:          | Mouse   |
| <b>Host Species:</b> | Goat  |

#### **Product Details**

| Background:   | Anti-Mouse IgG3 Peroxidase Antibody generated in goat detects reactivity to Mouse IgG3 (Gamma 3 chain). Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. IgG3 comprises almost 10% of the IgG subclass and has a high affinity for binding to the Fc receptor of phagocytic cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. |
|---------------|--|
| Synonyms:     | goat anti-MOUSE IgG3 Antibody peroxidase Conjugation, goat anti-MOUSE IgG3 HRP Conjugated Antibody   |
| Host Species: | Goat   |

|                      | Conjugated Antibody |
|----------------------|---------------------|
| <b>Host Species:</b> | Goat                |
| Specificity:         | IgG3                |
| Conjugate:           | Peroxidase (HRP)    |
| Clonality:           | Polyclonal          |
| Format:              | IgG                 |

# **Target Details**

| Reactivity:     | Mouse          |
|-----------------|----------------|
| Immunogen Type: | Native Protein |

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

| Immunogen:          | Mouse IgG3 heavy chain   |
|---------------------|--|
| Purity/Specificity: | Anti-MOUSE IgG3 (Gamma 3 chain) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-peroxidase, anti-Goat Serum, Mouse Serum and Mouse IgG3. No reaction was observed against Bovine, Human, and Rabbit Serum Proteins. Specificity was confirmed by ELISA at less than 1% of target signal. |

## **Application Details**

| <b>Tested Applications:</b> | ELISA  |
|-----------------------------|--|
| Suggested Applications:     | WB (Based on references)   |
| Application Note:           | Anti-Mouse IgG3 peroxidase antibody has been tested by ELISA and is ideal for western blotting, Immunohistochemistry and ELISA as well as other antibody detection methods and anti IgG2b antibody based assays. Mouse IgG3 secondary antibody is available in a variety of formats. HRP Conjugated Secondary Antibodies are ideal for western blot and are available in a variety of formats and conjugate types. |
| Assay Dilutions:            | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.  |
| ELISA:                      | 1:20,000 - 1:100,000   |
| IHC:                        | 1:1,000 - 1:5,000  |
| WB:                         | 1:2,000 - 1:10,000   |

### **Formulation**

| Physical State:        | Lyophilized  |
|------------------------|--|
| Concentration:         | 1.0 mg/mL by UV absorbance at 280 nm                                   |
| Buffer:                | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2             |
| Preservative:          | 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!               |
| Stabilizer:            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Reconstitution Volume: | 1.0 mL   |
| Reconstitution Buffer: | Restore with deionized water (or equivalent)                           |
|                        |  |

# **Shipping & Handling**

**Shipping Condition:** Ambient

www.rockland.com Page 2 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

| Storage Condition: | Store vial at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Anti-Mouse IgG HRP secondary antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
|--------------------|--|
| Expiration:        | Expiration date is one (1) year from date of receipt.  |

#### References

- Montero DA et al. A chimeric protein-based vaccine elicits a strong IgG antibody response and confers partial protection against Shiga toxin-producing Escherichia coli in mice. *Front Immunol.* (2023)
- Powers JM et al. Non-replicating adenovirus based Mayaro virus vaccine elicits protective immune responses and cross protects against other alphaviruses. *PLoS Negl Trop Dis.* (2021)
- Chen et al. Blocking fatty acid synthase inhibits tumor progression of human osteosarcoma by regulating the human epidermal growth factor receptor 2/phosphoinositide 3-kinase/protein kinase B signaling pathway in xenograft models. Experimental and Therapeutic Medicine (2017)
- Guo et al. Transcription factor NF②YA promotes a malignant phenotype by upregulating fatty acid synthase expression. Molecular Medicine Reports (2016)
- Secundino I et al. Salmonella porins induce a sustained, lifelong specific bactericidal antibody memory response. Immunology. (2006)

#### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3