



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for 610-152-043****Mouse IgG3 Antibody ATTO 488 Conjugated Pre-adsorbed****Overview**

<b>Description:</b>	Goat Anti-Mouse IgG3 (Gamma 3 chain) Antibody ATTO 488 Conjugated (Min X Bv, Hu, and Rb Serum Proteins) - 610-152-043
<b>Item No.:</b>	610-152-043
<b>Size:</b>	500 µg
<b>Applications:</b>	Dot Blot
<b>Reactivity:</b>	Mouse
<b>Host Species:</b>	Goat

**Product Details**

<b>Background:</b>	Anti-Mouse IgG3 ATTO 488 Antibody generated in goat detects reactivity to Mouse IgG3 (Gamma 3 chain). Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. IgG3 comprises almost 10% of the IgG subclass and has a high affinity for binding to the Fc receptor of phagocytic cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
<b>Synonyms:</b>	goat anti-MOUSE IgG3 Antibody ATTO488 Conjugation, goat anti-MOUSE IgG3 ATTO 488 Conjugated Antibody
<b>Host Species:</b>	Goat
<b>Specificity:</b>	IgG3
<b>Conjugate:</b>	ATTO 488
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG
<b>F/P Ratio:</b>	3.6

**Target Details**

<b>Reactivity:</b>	Mouse
--------------------	-------

<b>Immunogen Type:</b>	Native Protein
<b>Immunogen:</b>	Mouse IgG3 heavy chain
<b>Purity/Specificity:</b>	Anti-MOUSE IgG3 (Gamma 3 chain) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse Serum and Mouse IgG3. No reaction was observed against Bovine, Human, and Rabbit Serum Proteins. Specificity was confirmed by ELISA at less than 1% of target signal.

## Application Details

<b>Tested Applications:</b>	Dot Blot
<b>Application Note:</b>	Anti-Mouse IgG3 ATTO 488 Antibody has been tested by dot blot. Mouse IgG3 secondary antibody is available in a variety of formats. ATTO 488 Conjugated antibodies are designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. The emission spectra for this ATTO conjugate matches the principle output wavelengths of most common fluorescence instrumentation.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>FC:</b>	1:500 - 1:2,500
<b>FLISA:</b>	>1:20,000
<b>IF:</b>	>1:5,000
<b>WB:</b>	1:4,000 - 1:20,000


## Formulation

<b>Physical State:</b>	Lyophilized
<b>Concentration:</b>	1.0 mg/mL by UV absorbance at 280 nm
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
<b>Reconstitution Volume:</b>	500 µL
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)

## Shipping & Handling


<b>Shipping Condition:</b>	Ambient
<b>Storage Condition:</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Expiration:</b>	Expiration date is one (1) year from date of receipt.

## Images



### Dot Blot

Dot Blot of Goat anti-Mouse IgG3 Antibody ATTO 488  
 Conjugated. Antigen: Mouse IgG3. Load: Lane 1 - 50 ng Lane  
 2 - 16.67 ng Lane 3 - 5.56 ng Lane 4 - 1.85 ng Lane 5 - 0.62  
 ng. Primary antibody: none. Secondary antibody: Goat anti-  
 Mouse IgG3 Antibody ATTO 488 Conjugated at 1:1,000 for 60  
 min at RT. Block: MB-070 for 60 min at RT.



### Immunofluorescence Microscopy

ATTO® dyes can be used for multicolor immunofluorescent  
 detection with low background and high signal. Examples  
 shown are: A. Tubulin in PtK2- male Rat Kangaroo Kidney  
 Epithelial Cells was detected using ATTO 532 labeled  
 secondary antibody. B. Muscle alpha-actin was stained with  
 a mouse primary antibody and ATTO 488 anti-mouse IgG  
 (green) while Cytokeratin was stained with polyclonal rabbit  
 anti-cytokeratin and ATTO 647N anti-rabbit IgG (red). C.  
 HUVEC (Human umbilical vein endothelial cells) were stained  
 with anti- Vimentin-ATTO 532 (green), anti-E-Cadherin-ATTO  
 655 (red) and DAPI (blue). D. Rat colon sections were  
 stained with Anti-Aquaporin 3-ATTO 594 antibody. Hoechst  
 33342 (blue) is used as counterstain. Images provided  
 courtesy of Dr. Jörg Reichwein, ATTO-TEC GmbH

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.