



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 611-1502**Rabbit IgG (H&L) Antibody Alkaline Phosphatase Conjugated****Overview**

| | |
|----------------------|--|
| Description: | Goat Anti-Rabbit IgG (H&L) Antibody Alkaline Phosphatase Conjugated - 611-1502 |
| Item No.: | 611-1502 |
| Size: | 1 mg |
| Applications: | Dot Blot, ELISA, WB |
| Reactivity: | Rabbit |
| Host Species: | Goat |

Product Details

| | |
|----------------------|--|
| Background: | Anti-Rabbit IgG (H&L) Alkaline Phosphatase Antibody generated in goat detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. |
| Synonyms: | goat anti-Rabbit IgG Antibody Alkaline Phosphatase Conjugation, goat anti-Rabbit IgG Alk Phos Conjugated Antibody, goat anti-rabbit IgG heavy and light chain Antibody conjugated to Alkaline phosphatase |
| Host Species: | Goat |
| Specificity: | IgG (H&L) |
| Conjugate: | Alkaline Phosphatase (AP) |
| Clonality: | Polyclonal |
| Format: | IgG |

Target Details

| | |
|----------------------------|--|
| Reactivity: | Rabbit |
| Immunogen Type: | Native Protein |
| Immunogen: | Anti-Rabbit IgG (H&L) was produced by repeated immunization with rabbit whole IgG molecule in goat. |
| Purity/Specificity: | Anti-Rabbit IgG (H&L) Antibody Alkaline Phosphatase conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Goat Serum, Rabbit IgG and Rabbit Serum. |

Application Details

| | |
|-----------------------------|---|
| Tested Applications: | Dot Blot, ELISA, WB |
| Application Note: | Anti-Rabbit IgG (H&L) alkaline phosphatase conjugated antibody has been tested by ELISA, dot blot, and western blot and is suitable for immunoblotting (western or dot blot), ELISA, immunohistochemistry as well as other alkaline phosphatase-antibody based enzymatic assays requiring lot-to-lot consistency. |
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:3,000 - 1:25,000 |
| IHC: | 1:200 - 1:1,000 |
| WB: | 1:1,000 - 1:4,000 |


Formulation

| | |
|------------------------|---|
| Physical State: | Liquid (sterile filtered) |
| Concentration: | 1.0 mg/mL by UV absorbance at 280 nm |
| Buffer: | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 |
| Preservative: | 0.1% (w/v) Sodium Azide |
| Stabilizer: | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |

Shipping & Handling


| | |
|----------------------------|--|
| Shipping Condition: | Wet Ice |
| Storage Condition: | Store vial at 4° C before opening. DO NOT FREEZE. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. |
| Expiration: | Expiration date is one (1) year from date of receipt. |

Images




Figure

Typical experiment image where the complexes Ag–AbI–AbII are attached to the nanoparticles trapped in the magnetic traps and the fluorogenic substrate is flowed at different rates, where the measurement areas selected for the fluorescence analysis are shown. Measurement with a flow rate of (a) 1 µl/h, (b) 2 µl/h, (c) 5 µl/h, and (d) 10 µl/h. Experiment with anti-rabbit IgG labeled alkaline phosphatase was used (p/n 611-1502) and an anti-biotin rabbit IgG (p/n 100-4198) [AbI] concentration of 100 pg/ml. FIG. 6. PMID: 32038740.



Figure

Typical experiment image where the complexes Ag–AbI–AbII are attached to the nanoparticles trapped in the magnetic traps and the fluorogenic substrate is flowed at different rates, where the measurement areas selected for the fluorescence analysis are shown. Measurement with a flow rate of (a) 1 µl/h, (b) 2 µl/h, (c) 5 µl/h, and (d) 10 µl/h. Experiment with anti-rabbit IgG labeled alkaline phosphatase was used (p/n 611-1502) and an anti-biotin rabbit IgG (p/n 100-4198) [AbI] concentration of 100 pg/ml. FIG. 6. PMID: 32038740.




ELISA

ELISA results of Goat Anti-Rabbit IgG Antibody Alkaline Phosphatase Conjugated tested against purified Rabbit IgG protein. Each well was coated in duplicate with 10 µg of Rabbit IgG (p/n 011-0102). The starting dilution of antibody was 5 µg/ml and the X-axis represents the Log10 of a 3-fold dilution. The titer is 1:45,700. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% fish gelatin as blocking buffer and TMB substrate p/n TMBE-1000.



Dot Blot

Dot Blot of Rabbit IgG Antibody Alkaline Phosphatase Conjugated. Antigen: Rabbit IgG. Load: Lane 1 - 200ng, Lane 2 - 66.7ng, Lane 3 - 22.2ng, Lane 4 - 7.4ng, Lane 5 - 2.5ng. Primary antibody: none. Secondary antibody: Rabbit IgG Antibody Alkaline Phosphatase Conjugated at 1:1,000 for 60 min at RT. Block: MB-070 for 60 min at RT. Reaction visualized using alkaline phosphatase substrate for 30 seconds at RT.



Western Blot

Western Blot of Goat anti-Rabbit IgG Antibody Alkaline Phosphatase Conjugated. Lane 1: Rabbit IgG. Lane 2: None. Load: 100 ng per lane. Primary Antibody: None. Secondary antibody: Alkaline Phosphatase goat secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 55 and 28 kDa, 55 kDa for Rabbit IgG. Other Band(s): None.

References

- Guevara-Pantoja PE et al. Micro–nanoparticles magnetic trap: Toward high sensitivity and rapid microfluidic continuous flow enzyme immunoassay. *Biomicrofluidics*. (2020)
- Mueller A et al. In vitro assembly of Tobacco mosaic virus coat protein variants derived from fission yeast expression clones or plants. *J Virol Methods*. (2010)
- Copello JA et al. Lack of effect of cADP-ribose and NAADP on the activity of skeletal muscle and heart ryanodine receptors. *Cell Calcium*. (2001)
- DeLamatre JG et al. Influence of dietary fat on the effect of endotoxin on murine hepatic peroxisomes. *Hepatology*. (1996)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.