

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 618-100-007

# Ferret IgM (mu chain) Antibody Rhodamine Conjugated

#### **Overview**

| Description:  | Goat Anti-Ferret IgM (mu chain) Antibody Rhodamine Conjugated - 618-100-007 |  |  |  |
|---------------|---|--|--|--|
| Item No.:     | 618-100-007   |  |  |  |
| Size:         | 1 mg  |  |  |  |
| Reactivity:   | Ferret  |  |  |  |
| Host Species: | Goat  |  |  |  |

### **Product Details**

| Background:   | Anti-Ferret IgM antibody specifically detects ferret IgM. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum. Anti-Ferret IgM antibody is ideal for investigators in Immunology, Microbiology, and Cell Biology. |
|---------------|--|
| Synonyms:     | goat anti-Ferret IgM mu chain Antibody rhodamine conjugation, goat anti-ferret IgM mu TRITC conjugated antibody  |
| Host Species: | Goat   |
| Specificity:  | IgM μ chain  |

Rhodamine (TRITC)

Polyclonal

IgG

### **Target Details**

Conjugate:

Clonality: Format:

| Reactivity: | Ferret                    |  |  |  |
|-------------|---------------------------|--|--|--|
| Immunogen:  | Ferret IgM whole molecule |  |  |  |

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

**Purity/Specificity:** This product was prepared from monospecific antiserum by immunoaffinity chromatography

using Ferret IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Ferret IgM and Ferret Serum. No reaction was observed against other

ferret heavy or light chain proteins.

#### **Application Details**

| Application Note: | This product is designed for immunofluorescence microscopy, fluorescence based plate assays     |
|-------------------|---|
|                   | (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, |

including multicolor imaging, utilizing various commercial platforms.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be

listed below.

**IF:** 1:500-1:2,500

#### **Formulation**

| Physical State: | Lyophilized |
|-----------------|-------------|
|-----------------|-------------|

| Concentration: | 1 mg | /mL l | by U\ | / absorbance at | 280 nm |
|----------------|------|-------|-------|-----------------|--------|
|----------------|------|-------|-------|-----------------|--------|

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Preservative:** 0.01% (w/v) Sodium Azide

Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

**Reconstitution Volume:** 1.0 mL

**Reconstitution Buffer:** Restore with deionized water (or equivalent)

## **Shipping & Handling**

| ς | hi | nni | ing | Cond  | iti | ion:  | Α             | m | hi  | en  | ıt |
|---|----|-----|-----|-------|-----|-------|---------------|---|-----|-----|----|
| _ |    | יאא |     | COIIG |     | 1011. | $\overline{}$ |   | O I | CII | ·  |

**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° Control of the vision and the vision of freeze at -20° Control of the vision of

C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use

undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

#### **Disclaimer**

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3