



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 706-105-002

F(ab')₂ Guinea Pig IgG (H&L) Antibody Alkaline Phosphatase Conjugated

Overview

| | |
|----------------------|---|
| Description: | Goat F(ab') ₂ Anti-Guinea Pig IgG (H&L) Antibody Alkaline Phosphatase Conjugated - 706-105-002 |
| Item No.: | 706-105-002 |
| Size: | 500 µg |
| Reactivity: | Guinea Pig |
| Host Species: | Goat |

Product Details

| | |
|----------------------|---|
| Background: | F(ab') ₂ Anti-Guinea Pig IgG Alkaline Phosphatase antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab') ₂ fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab') ₂ fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab') ₂ fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab') ₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays. |
| Synonyms: | Goat F(ab') ₂ Anti-Guinea Pig IgG Antibody Alkaline Phosphatase Conjugation, Goat Fab2 Anti-Guinea Pig IgG alk phos Conjugated Antibody |
| Host Species: | Goat |
| Specificity: | IgG (H&L) |
| Conjugate: | Alkaline Phosphatase (AP) |
| Clonality: | Polyclonal |
| Format: | IgG F(ab') ₂ |

Target Details

| | |
|--------------------|-------------------------------|
| Reactivity: | Guinea Pig |
| Immunogen: | Guinea Pig IgG whole molecule |

| | |
|----------------------------|---|
| Purity/Specificity: | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c). |
|----------------------------|---|

Application Details

| | |
|--------------------------|--|
| Application Note: | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 µg of Guinea pig IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product. |
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:2,000 - 1:10,000 |
| IHC: | 1:200 - 1:1,000 |
| WB: | 1:500 - 1:2,500 |

Formulation

| | |
|------------------------|---|
| Physical State: | Liquid (sterile filtered) |
| Concentration: | 1.0 mg/mL by UV absorbance at 280 nm |
| Buffer: | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 |

Shipping & Handling

| | |
|----------------------------|--|
| Shipping Condition: | Wet Ice |
| Storage Condition: | Store vial at 4° C before opening. DO NOT FREEZE. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. |
| Expiration: | Expiration date is one (1) year from date of receipt. |

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.