



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for 711-1122****F(ab')<sub>2</sub> Rabbit IgG (H&L) Antibody Pre-Adsorbed****Overview**

<b>Description:</b>	Goat F(ab') <sub>2</sub> Anti-Rabbit IgG (H&L) Antibody (Min X Bv Ch Gt GP Ham Hs Hu Ms Rt & Sh Serum Proteins) - 711-1122
<b>Item No.:</b>	711-1122
<b>Size:</b>	500 µg
<b>Applications:</b>	WB
<b>Reactivity:</b>	Rabbit
<b>Host Species:</b>	Goat

**Product Details**

<b>Background:</b>	F(ab') <sub>2</sub> Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab) <sub>2</sub> fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab) <sub>2</sub> fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab) <sub>2</sub> fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab') <sub>2</sub> Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.
<b>Synonyms:</b>	Goat F(ab') <sub>2</sub> Anti-Rabbit IgG Antibody Pre-Adsorbed, Goat Fab2 Anti-Rabbit IgG Antibody
<b>Host Species:</b>	Goat
<b>Specificity:</b>	IgG (H&L)
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG F(ab') <sub>2</sub>

**Target Details**

<b>Reactivity:</b>	Rabbit
<b>Immunogen:</b>	Rabbit IgG whole molecule

<b>Purity/Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rat and Sheep Serum Proteins.
----------------------------	--

## Application Details

<b>Suggested Applications:</b>	WB (Based on references)
<b>Application Note:</b>	F(ab') <sub>2</sub> Anti-Rabbit IgG Antibody is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:20,000
<b>IHC:</b>	1:1,000 - 1:5,000
<b>WB:</b>	1:2,000 - 1:10,000

## Formulation


<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	1.0 mg/mL by UV absorbance at 280 nm
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	None

## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
<b>Storage Condition:</b>	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images




### Western Blot

KIF20B shRNA efficiency. (B) The shRNA efficiency measured by western blotting. Primary antibodies used rabbit anti-KIF20B antibody [1:100] and mouse anti-β-actin [1:500] with goat anti-rabbit [1:10,000] p/n 711–1122 and goat anti-mouse Fc secondary antibody [1:10,000] p/n 110–1103. \*P<0.05 vs. control. KIF, kinesin family member; shRNA, short hairpin RNA. Figure 2. PMID: 30664160.

### Western Blot

Western Blot results using Goat F(ab')<sub>2</sub> Anti-Rabbit IgG. Effects of RAB38 silencing on the proliferation ability of PANC-1 and BxPC-3 cells. The expression levels of the proliferation-associated proteins (C) Ki67 and (D) PCNA in PANC-1 and BxPC-3 cells were detected by western blot analysis. Data are reported as the mean ± standard deviation (n=3). \*P<0.05 vs. control group. shRNA, small hairpin RNA; RAB38, Ras-related GTP-binding protein 38; OD, optical density; PCNA, proliferating cell nuclear antigen. Figure 3. PMID: 30569114.



#### SDS-PAGE

SDS-PAGE results of Goat F(ab')<sub>2</sub> Anti-Rabbit IgG (H&L) Antibody. Lane 1: reduced F(ab')<sub>2</sub> anti-Mouse (lot# 36997). Lane 2: reduced F(ab')<sub>2</sub> anti-Rabbit (lot# 36998). Lane 3: Opal Prestained Ladder - MB-210-0500. Lane 4: non-reduced F(ab')<sub>2</sub> anti-Mouse. Lane 5: non-reduced F(ab')<sub>2</sub> anti-Rabbit. Load: 1.0ug. 4-20% Lonza SDS-PAGE; Coomassie Stained; BioRad ChemiDoc Imaged.

## References

- Uchida Y et al. Generation of antagonistic monoclonal antibodies against the neoepitope of active mouse interleukin (IL)-18 cleaved by inflammatory caspases. *Arch Biochem Biophys.* (2022)
- Li et al. Kinesin family member 20B regulates tongue cancer progression by promoting cell proliferation. *Molecular Medicine Reports* (2019)
- Li et al. High expression of RAB38 promotes malignant progression of pancreatic cancer. *Molecular Medicine Reports* (2018)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.