

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 712-1122

F(ab')2 Rat IgG Fc Antibody Pre-Adsorbed

Overview

| Description: | Goat F(ab')2 Anti-Rat IgG Fc Antibody (Min X Bv Hs & Hu Serum Proteins) - 712-1122 |
|---------------|--|
| Item No.: | 712-1122 |
| Size: | 1 mg |
| Reactivity: | Rat |
| Host Species: | Goat |

Product Details

| Background: | F(ab')2 Anti-Rat IgG F(c) Antibody was generated in goat and detects specifically Rat IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. |
|----------------------|---|
| Synonyms: | Goat F(ab')2 Anti-Rat IgG F(c) Antibody, Goat Fab2 Anti-Rat IgG Fc Antibody |
| Host Species: | Goat |
| Specificity: | IgG Fc |
| Clonality: | Polyclonal |
| Format: | IgG F(ab')2 |

Target Details

| Reactivity: | Rat |
|---------------------|---|
| Immunogen Type: | Native Protein |
| Immunogen: | Anti-Rat IgG F(c) was produced by repeated immunization with Rat F(c) fragment in goat. |
| Purity/Specificity: | Anti-Rat F(c) antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG, Rat IgG F(c) and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(ab) or Bovine, Horse and Human Serum Proteins. |

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

Application Details

| Application Note: | F(ab')2 Anti-Rat IgG F(c) Antibody is suitable for immunoblotting, western blot, dot blot, ELISA, and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency. |
|-------------------|--|
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:10,000 - 1:50,000 |
| IHC: | 1:1,000 - 1:3,000 |
| WB: | 1:2,000 - 1:5,000 |

Formulation

| Physical State: | Liquid (sterile filtered) |
|-----------------|--|
| Concentration: | 1.0 mg/mL by UV absorbance at 280 nm |
| Buffer: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Preservative: | 0.01% (w/v) Sodium Azide |
| Stabilizer: | None |

Shipping & Handling

| Shipping Condition: | Wet Ice |
|---------------------|--|
| Storage Condition: | Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. |
| Expiration: | Expiration date is one (1) year from date of receipt. |


Images

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823



SDS-PAGE

SDS-PAGE results of Goat F(ab')2 Anti-Rat IgG F(c) Antibody. Lane 1: reduced Goat F(ab')2 anti-Rat. Lane 2: Opal Prestained Ladder - MB-210-0500. Lane 3: non-reduced Goat F(ab')2 anti-Rat. Load: 1.0ug. 4-20% Lonza SDS-PAGE; Coomassie Stained; BioRad ChemiDoc Imaged.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3