



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



**Datasheet for 712-1131****F(ab')<sub>2</sub> Rat IgG F(ab')<sub>2</sub> Antibody Pre-Adsorbed****Overview**

|                      |   |
|----------------------|---|
| <b>Description:</b>  | Goat F(ab') <sub>2</sub> Anti-Rat IgG F(ab') <sub>2</sub> Antibody (Min X Bv Hs & Hu Serum Proteins) - 712-1131 |
| <b>Item No.:</b>     | 712-1131  |
| <b>Size:</b>         | 1 mg  |
| <b>Reactivity:</b>   | Rat   |
| <b>Host Species:</b> | Goat  |

**Product Details**

|                      |  |
|----------------------|--|
| <b>Background:</b>   | F(ab') <sub>2</sub> Anti-Rat IgG F(ab') <sub>2</sub> Antibody generated in goat detects Rat F(ab') <sub>2</sub> . Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab') <sub>2</sub> Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays. |
| <b>Synonyms:</b>     | Goat F(ab') <sub>2</sub> Anti-Rat IgG F(ab') <sub>2</sub> Antibody, Goat Fab2 Anti-Rat IgG Fab2 Antibody   |
| <b>Host Species:</b> | Goat   |
| <b>Specificity:</b>  | IgG F(ab') <sub>2</sub>  |
| <b>Clonality:</b>    | Polyclonal   |
| <b>Format:</b>       | IgG F(ab') <sub>2</sub>  |

**Target Details**

|                    |                                      |
|--------------------|--------------------------------------|
| <b>Reactivity:</b> | Rat                                  |
| <b>Immunogen:</b>  | Rat IgG F(ab') <sub>2</sub> fragment |

|                            |  |
|----------------------------|--|
| <b>Purity/Specificity:</b> | F(ab') <sub>2</sub> Rat IgG F(ab') <sub>2</sub> Antibody Pre-Adsorbed was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG, Rat IgG F(ab') <sub>2</sub> and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(c) or Bovine, Horse and Human Serum Proteins. |
|----------------------------|--|

## Application Details

|                          |   |
|--------------------------|---|
| <b>Application Note:</b> | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications. |
|--------------------------|---|

|                         |   |
|-------------------------|---|
| <b>Assay Dilutions:</b> | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| <b>ELISA:</b>           | 1:20,000 - 1:100,000  |
| <b>IHC:</b>             | 1:1,000 - 1:5,000   |
| <b>WB:</b>              | 1:2,000 - 1:10,000  |

## Formulation

|                        |   |
|------------------------|---|
| <b>Physical State:</b> | Liquid (sterile filtered)                               |
| <b>Concentration:</b>  | 1.3 mg/mL by UV absorbance at 280 nm                    |
| <b>Buffer:</b>         | 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2 |
| <b>Preservative:</b>   | None  |
| <b>Stabilizer:</b>     | None  |

## Shipping & Handling

|                            |  |
|----------------------------|--|
| <b>Shipping Condition:</b> | Wet Ice  |
| <b>Storage Condition:</b>  | Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. |
| <b>Expiration:</b>         | Expiration date is one (1) year from date of receipt.  |

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.