

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

### Datasheet for 813-4102 Fab Sheep IgG (H&L) Antibody

### **Overview**

Description:	Rabbit Fab Anti-Sheep IgG (H&L) Antibody - 813-4102
Item No.:	813-4102
Size:	500 µg
Reactivity:	Sheep
Host Species:	Rabbit

### **Product Details**

Background:	Fab Anti-Sheep IgG (H&L) Antibody generated in rabbit detects immunoglobulin g from sheep, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross- reactivity, and host-species source and fragment composition.
Synonyms:	Rabbit Fab Anti-Sheep IgG Antibody, Rabbit Fab Fragment Anti-Sheep IgG Antibody
Host Species:	Rabbit
Specificity:	IgG (H&L)
Clonality:	Polyclonal
Format:	lgG Fab

### **Target Details**

Reactivity:	Sheep
Immunogen:	Sheep IgG whole molecule



www.rockland.com tech@rockland.com +1 484.791.3823

# Purity/Specificity:This product was prepared from monospecific antiserum by immunoaffinity chromatography<br/>using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any<br/>unwanted reactivities, papain digestion and chromatographic separation. Assay by<br/>immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. No<br/>reaction was observed against anti-Papain or anti-Rabbit IgG F(c).

### **Application Details**

Application Note:	Suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:100,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000

### **Formulation**

Physical State:	Liquid (sterile filtered)
Concentration:	0.5 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

### **Shipping & Handling**

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

### Disclaimer



#### Order online now!

www.rockland.com tech@rockland.com +1 484.791.3823

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.