



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for MB-006****10X TE pH 7.5****Overview**

<b>Description:</b>	10X TE pH 7.5 (100 mM Tris HCl 10 mM EDTA) DEPC Treated - MB-006
<b>Item No.:</b>	MB-006
<b>Size:</b>	1 L
<b>Applications:</b>	IHC

**Product Details**

<b>Background:</b>	TE buffer is a commonly used buffer solution in molecular biology, especially in procedures involving DNA, cDNA or RNA. "TE" is derived from its components: Tris, a common pH buffer, and EDTA, a molecule that chelates cations like Mg <sup>2+</sup> . The purpose of TE buffer is to solubilize DNA or RNA, while protecting it from degradation.
<b>Synonyms:</b>	10X TE pH 7.5 (100 mM Tris HCl 10 mM EDTA) Solution, 10X TE pH 7.5 DEPC TREATED Solution

**Target Details**

<b>Purity/Specificity:</b>	This product was aseptically filtered through a Millipore 0.22 micron filter into clean, pre-sterilized containers. The product was tested on trypticase soy agar for 24 hours, 48 hours and 72 hours and was found to be negative for bacteria.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li><a href="#">MB-006 SDS</a></li></ul>

**Application Details**

<b>Suggested Applications:</b>	IHC (Based on references)
<b>Application Note:</b>	This product is a concentrated stock solution and should be diluted appropriately with distilled, deionized water or equivalent to its final working concentration. This buffer consists of 100 mM Tris HCl and 10 mM EDTA at a pH of 7.5. Meticulously prepared using ultra pure reagents dissolved in DEPC treated molecular biology grade water.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.


## Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	10X
Buffer:	See application note.
Preservative:	None
Stabilizer:	None

## Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store container at room temperature (18° to 26° C) prior to opening. If desired, the solution may be stored at 4° C or less. Some salts may precipitate out of solution at lower temperature. Allow buffer to equilibrate to room temperature (18° to 26° C) to restore solubility of some salts.
Expiration:	Expiration date is six (6) months from date of receipt.

## Images



### Immunofluorescence Microscopy

Immunohistochemistry for MLH1 and MSH2 protein expression (hematoxylin, 400x). (a) Normal colorectal tissue with no antibody staining for MLH1 (negative control). (b) Normal colorectal tissue with MLH1 antibody staining (positive control). (c) Tumor with absence of MLH1 protein expression. (d) Tumor with MLH1 protein expression. (e) Normal colorectal tissue with no antibody staining for MSH2 (negative control). (f) Normal colorectal tissue with MSH2 antibody staining (positive control). (g) Tumor with absence of MSH2 protein expression. (h) Tumor with MSH2 protein expression. The slides underwent antigen retrieval with 10X Tris-ethylenediamine tetraacetic acid buffer, pH 7.5 (p/n MB-006) for 45 minutes on boiling water. Fig 2. PMID: 20012372.

**Bottle**

10X TE pH 7.5 (100 mM Tris HCl 10 mM EDTA) DEPC Treated

## References

- De Jesus-Monge, WE. et al. Mismatch repair protein expression and colorectal cancer in Hispanics from Puerto Rico. *Familial Cancer* (2010)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.