



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



EFP (h2): 293T Lysate: sc-115553

BACKGROUND

EFP (estrogen-responsive finger protein) is a transcription factor, the content of which is regulated by estrogen. It has been identified as a member of the RING-finger family, a family of proteins containing a Zn²⁺ binding domain designated the C3HC4 or RING finger. EFP also contains two B box domains and a coiled-coil region (a transactivation domain), which are characteristic of a subgroup of the RING-finger family. Estrogen regulates the growth, differentiation and function of target cells in a variety of tissues; however, few genes have been shown to be directly regulated by estrogen. It has been speculated that EFP may mediate estrogen activity in a signaling cascade in which estrogen-ER binding to the estrogen responsive element (ERE) downstream of the EFP gene upregulates EFP gene expression. The EFP gene product may then activate transcription of secondary estrogen responsive genes. Additional studies indicate that the EFP promoter may be regulated by multiple elements and their interacting factors.

REFERENCES

1. Evans, R.M. 1988. The steroid and thyroid hormone receptor superfamily. *Science* 240: 889-895.
2. Green, S., et al. 1988. Nuclear receptors enhance our understanding of transcription regulation. *Trends Genet.* 4: 309-314.
3. Inoue, S., et al. 1993. Genomic binding-site cloning reveals an estrogen-responsive gene that encodes a RING-finger protein. *Proc. Natl. Acad. Sci. USA* 90: 11117-11121.
4. Orimo, A., et al. 1995. Molecular cloning, structure, and expression of mouse estrogen-responsive finger protein EFP. Co-localization with estrogen receptor mRNA in target organs. *J. Biol. Chem.* 270: 24406-24013.
5. Borden, K.L., et al. 1995. The solution structure of the RING-finger domain from the acute promyelocytic leukaemia proto-oncoprotein PML. *EMBO J.* 14: 1532-1541.
6. Ikeda, K., et al. 1997. Multiple regulatory elements and binding proteins of the 5'-flanking region of the human estrogen-responsive finger protein (EFP) gene. *Biochem. Biophys. Res. Commun.* 236: 765-771.

CHROMOSOMAL LOCATION

Genetic locus: TRIM25 (human) mapping to 17q22.

PRODUCT

EFP (h2): 293T Lysate represents a lysate of human EFP transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

EFP (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive EFP antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.