



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PYK2 (h): 293T Lysate: sc-115595

BACKGROUND

Focal adhesion kinase (FAK) was initially identified as a substrate for the intrinsic protein tyrosine kinase activity of Src-encoded pp60. The deduced amino acid sequence of FAK p125 has shown it to be a cytoplasmic protein tyrosine kinase whose sequence and structural organization are unique compared to other protein families described. A putative new member of the FAK family, designated PYK2 (proline-rich tyrosine kinase 2), exhibits 61% sequence identity with FAK over its kinase domain. PYK2 (also designated CAK β or RAFTK) is highly expressed in the central nervous system. Activation of the kinase leads to modulation of ion channel function and the activation of the MAPK signaling pathway. PYK2 is rapidly phosphorylated on tyrosine residues in response to stimuli that increase intracellular calcium levels and compounds that activate members of the PKC family of kinases, such as phorbol esters.

REFERENCES

- Schaller, M.D., Borgman, C.A., Cobb, B.S., Vines, R.R., Reynolds, A.B. and Parsons, J.T. 1992. pp125FAK, a structurally distinctive protein-tyrosine kinase associated with focal adhesions. *Proc. Natl. Acad. Sci. USA* 89: 5192-5196.
- Lipfert, L., Haimovich, B., Schaller, M.D., Cobb, B.S., Parsons, J.T. and Brugge, J.S. 1992. Integrin-dependent phosphorylation of the protein tyrosine kinase pp125FAK in platelets. *J. Cell Biol.* 119: 905-912.
- Hanks, S.K., Calallo, M.B., Harper, M.C. and Patel, S.K. 1992. Focal adhesion protein-tyrosine kinase phosphorylated in response to cell attachment to Fibronectin. *Proc. Natl. Acad. Sci. USA* 89: 8487-8491.
- Guan, J.L. and Shalloway, D. 1992. Regulation of focal adhesion-associated protein tyrosine kinase by both cellular adhesion and oncogenic transformation. *Nature* 359: 690-692.
- Schaller, M.D., Hildebrand, J.D., Shannon, J.D., Fox, J.W., Vines, R.R. and Parsons, J.T. 1994. Autophosphorylation of the focal adhesion-associated protein tyrosine kinase, pp125FAK, directs SH2-dependent binding of pp60Src. *Mol. Cell. Biol.* 14: 1680-1688.
- Lev, S., Moreno, H., Martinez, R., Canoll, P., Peles, E., Musacchio, J.M., Plowman, G.D., Rudy, B. and Schlessinger, J. 1995. Protein tyrosine kinase PYK2 involved in Ca²⁺-induced regulation of ion channel and MAP kinase functions. *Nature* 376: 737-745.

CHROMOSOMAL LOCATION

Genetic locus: PTK2B (human) mapping to 8p21.2.

PRODUCT

PYK2 (h): 293T Lysate represents a lysate of human PYK2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

PYK2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive PYK2 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PYK2 (E-3): sc-393181 is recommended as a positive control antibody for Western Blot analysis of enhanced human PYK2 expression in PYK2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PYK2 (E-3): sc-393181. Near-infrared western blot analysis of PYK2 expression in non-transfected: sc-117752 (A) and human PYK2 transfected: sc-115595 (B) 293T whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgG κ BP-CFL 790: sc-516181.



PYK2 (E-3): sc-393181. Near-infrared western blot analysis of PYK2 expression in non-transfected: sc-117752 (A) and human PYK2 transfected: sc-115595 (B) 293T whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgG κ BP-CFL 680: sc-516180.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.