



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NP2 (h2): 293T Lysate: sc-115612

BACKGROUND

Long pentraxins are a family of highly conserved proteins that are expressed in the brain and central nervous system, and form multimeric complexes. Neuronal pentraxin 1 (NP1), NP2, and neuronal pentraxin receptor (NPR) are members of the long pentraxins that represent a neuronal uptake pathway that may function during synapse formation and remodeling. The NP1 gene is located on chromosome 17q25.3 and the protein product mediates the uptake of synaptic material, including the presynaptic snake venom toxin, taipoxin. NP2, whose function is unknown, is located on chromosome 7q22.1 and like NP1 contains several potential N-linked glycosylation sites. NPR is expressed on the cell membrane and can form heteropentamers with NP1 and NP2 that can be released from the cell membrane by proteolysis.

REFERENCES

1. Hsu, Y.C. and Perin, M.S. 1995. Human neuronal pentraxin II (NPTX2): conservation, genomic structure, and chromosomal localization. *Genomics* 28: 220-227.
2. Dodds, D.C., Omeis, I.A., Cushman, S.J., Helms, J.A. and Perin, M.S. 1997. Neuronal pentraxin receptor, a novel putative integral membrane pentraxin that interacts with neuronal pentraxin 1 and 2 and taipoxin-associated calcium-binding protein 49. *J. Biol. Chem.* 272: 21488-21494.
3. Kirkpatrick, L.L., Matzuk, M.M., Dodds, D.C. and Perin, M.S. 2000. Biochemical interactions of the neuronal pentraxins. Neuronal pentraxin (NP) receptor binds to taipoxin and taipoxin-associated calcium-binding protein 49 via NP1 and NP2. *J. Biol. Chem.* 275: 17786-17792.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 600750. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Park, J.K., Ryu, J.K., Lee, K.H., Lee, J.K., Yoon, W.J., Lee, S.H., Yoo, J.W., Woo, S.M., Lee, G.Y., Lee, C.H., Kim, Y.T. and Yoon, Y.B. 2007. Quantitative analysis of NPTX2 hypermethylation is a promising molecular diagnostic marker for pancreatic cancer. *Pancreas* 35: 9-15.
6. Marui, T., Koishi, S., Funatogawa, I., Yamamoto, K., Matsumoto, H., Hashimoto, O., Ishijima, M., Nanba, E., Nishida, H., Sugiyama, T., Kasai, K., Watanabe, K., Kano, Y., Kato, N. and Sasaki, T. 2007. No association between the neuronal pentraxin II gene polymorphism and autism. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 31: 940-943.
7. Moran, L.B., Hickey, L., Michael, G.J., Derkacs, M., Christian, L.M., Kalaitzakis, M.E., Pearce, R.K. and Graeber, M.B. 2008. Neuronal pentraxin II is highly upregulated in Parkinson's disease and a novel component of Lewy bodies. *Acta Neuropathol.* 115: 471-478.

CHROMOSOMAL LOCATION

Genetic locus: NPTX2 (human) mapping to 7q22.1.

PRODUCT

NP2 (h2): 293T Lysate represents a lysate of human NP2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

NP2 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive NP2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

NP2 (G-9): sc-166035 is recommended as a positive control antibody for Western Blot analysis of enhanced human NP2 expression in NP2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



NP2 (G-9): sc-166035. Western blot analysis of NP2 expression in non-transfected: sc-117752 (A) and human NP2 transfected: sc-115612 (B) 293T whole cell lysates.



NP2 (G-2): sc-365759. Western blot analysis of NP2 expression in non-transfected: sc-117752 (A) and human NP2 transfected: sc-115612 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.