



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cosmc (h): 293T Lysate: sc-116086

BACKGROUND

Cosmc, or C1GALT1-specific chaperone 1, is a 318 amino acid protein encoded by the human gene C1GALT1C1. Cosmc is believed to be a chaperone required for the generation of 1 O-glycan Gal- β 1-3GalNAc- α 1-Ser/Thr (Tn antigen), which is a precursor for many extended O-glycans in glycoproteins. Cosmc may also function as a specific molecular chaperone assisting the folding/stability of core 1 β -3-galactosyltransferase (C1GALT1). Cosmc is a single-pass type II membrane protein. Cosmc is ubiquitously expressed in all tissue types. It is most abundantly expressed in small intestine, stomach, salivary gland, kidney and testis, and at intermediate levels in whole brain, cerebellum, spinal cord, thymus, spleen, trachea, lung, pancreas, ovary and uterus. Defects in C1GALT1C1 are the cause of Tn syndrome, a rare autoimmune disease caused by somatic mutation in the C1GALT1C1 gene in which subpopulations of blood cells of all lineages carry an incompletely glycosylated Tn antigen, effecting red cells and platelets and leading to anemia, leukopenia and thrombocytopenia. Tn-polyagglutinability is sometimes associated with leukemia or is a preleukemic state.

REFERENCES

1. Zhang, Q.H., Ye, M., Wu, X.Y., Ren, S.X., Zhao, M., Zhao, C.J., Fu, G., Shen, Y., Fan, H.Y., Lu, G., Zhong, M., Xu, X.R., Han, Z.G., Zhang, J.W., Tao, J., Huang, Q.H., Zhou, J., Hu, G.X., Gu, J., Chen, S.J. and Chen, Z. 2000. Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34⁺ hematopoietic stem/progenitor cells. *Genome Res.* 10: 1546-1560.
2. Kudo, T., Iwai, T., Kubota, T., Iwasaki, H., Takayama, Y., Hiruma, T., Inaba, N., Zhang, Y., Gotoh, M., Togayachi, A. and Narimatsu, H. 2002. Molecular cloning and characterization of a novel UDP-Gal:GalNAc(a) peptide β 1,3-galactosyltransferase (C1Gal-T2), an enzyme synthesizing a core 1 structure of O-glycan. *J. Biol. Chem.* 277: 47724-47731.
3. Ju, T. and Cummings, R.D. 2002. A unique molecular chaperone Cosmc required for activity of the mammalian core 1 β 3-galactosyltransferase. *Proc. Natl. Acad. Sci. USA* 99: 16613-16618.
4. Ju, T. and Cummings, R.D. 2005. Protein glycosylation: chaperone mutation in Tn syndrome. *Nature* 437: 1252-1252.
5. Qin, W., Zhou, Q., Yang, L.C., Li, Z., Su, B.H., Luo, H. and Fan, J.M. 2005. Peripheral B lymphocyte β 1,3-galactosyltransferase and chaperone expression in immunoglobulin A nephropathy. *J. Intern. Med.* 258: 467-477.
6. Ju, T., Zheng, Q. and Cummings, R.D. 2006. Identification of core 1 O-glycan T-synthase from *Caenorhabditis elegans*. *Glycobiology* 16: 947-958.
7. Schietinger, A., Philip, M., Yoshida, B.A., Azadi, P., Liu, H., Meredith, S.C. and Schreiber, H. 2006. A mutant chaperone converts a wildtype protein into a tumor-specific antigen. *Science* 314: 304-308.
8. LocusLink Report (LocusID: 29071). <http://www.ncbi.nlm.nih.gov/LocusLink/>

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: C1GALT1C1 (human) mapping to Xq24.

PRODUCT

Cosmc (h): 293T Lysate represents a lysate of human Cosmc transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Cosmc (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Cosmc antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.