



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NFκB p50 (h): 293T Lysate: sc-116112

BACKGROUND

Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NFκB (p50 and p65) and the *Drosophila* maternal morphogen, dorsal. These proteins share sequence homology over a region of 300 amino acids at their NH₂-terminus, the region that contains their DNA binding and dimerization domains. The DNA binding activity of NFκB is activated and rapidly transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins have been described. These proteins, designated p105 and p100, are highly related but map on different chromosomes. The p105 (p110) precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated PDI, binds to p50 and regulates its activity.

REFERENCES

1. Meyer, R., et al. 1991. Cloning of the DNA-binding subunit of human nuclear factor κB: the level of its mRNA is strongly regulated by phorbol ester or tumor necrosis factor α. *Proc. Natl. Acad. Sci. USA* 88: 966-970.
2. Schmid, R.M., et al. 1991. Cloning of an NFκB subunit which stimulates HIV transcription in synergy with p65. *Nature* 352: 733-736.
3. Ballard, D.W., et al. 1992. The 65 kDa subunit of human NFκB functions as a potent transcriptional activator and a target for v-Rel-mediated repression. *Proc. Natl. Acad. Sci. USA* 89: 1875-1879.
4. Hatada, E.N., et al. 1992. The Ankyrin repeat domains of the NFκB precursor p105 and the proto-oncogene Bcl-3 act as specific inhibitors of NFκB DNA binding. *Proc. Natl. Acad. Sci. USA* 89: 2489-2493.
5. Henkel, T., et al. 1992. Intramolecular masking of the nuclear location signal and dimerization domain in the precursor for the p50 NFκB subunit. *Cell* 69: 1121-1133.
6. Beg, A.A., et al. 1995. Embryonic lethality and liver degeneration in mice lacking the RelA component of NFκB. *Nature* 376: 167-170.
7. Huxford, T., et al. 1998. The crystal structure of the IκBα/NFκB complex reveals mechanisms of NFκB inactivation. *Cell* 95: 759-770.
8. Chen, L.F., et al. 2001. Duration of nuclear NFκB action regulated by reversible acetylation. *Science* 293: 1653-1657.
9. Zhang, J., et al. 2007. NFκB1/p50 is not required for tumor necrosis factor-stimulated growth of primary mammary epithelial cells: implications for NFκB2/p52 and RelB. *Endocrinology* 148: 268-278.

CHROMOSOMAL LOCATION

Genetic locus: NFKB1 (human) mapping to 4q24.

PRODUCT

NFκB1 p50 (h): 293T Lysate represents a lysate of human NFκB1 p50 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.


APPLICATIONS

NFκB1 p50 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive NFκB1 p50 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

NFκB p50 (4D1): sc-53744 is recommended as a positive control antibody for Western Blot analysis of enhanced human NFκB1 p50 expression in NFκB1 p50 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



NFκB p50 (4D1): sc-53744. Western blot analysis of NFκB p50 expression in non-transfected: sc-117752 (A) and human NFκB p50 transfected: sc-116112 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.