



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



apoL-VI (h2): 293T Lysate: sc-116232

BACKGROUND

The apolipoprotein L gene family maps to a region on chromosome 22 and encodes six highly homologous proteins, designated apoL-I, apoL-II, apoL-III, apoL-IV, apoL-V and apoL-VI, all of which function as components of plasma lipoproteins. ApoL-VI (Apolipoprotein L-VI), also known as APOL6, is a 343 amino acid protein that localizes to the cytoplasm and belongs to the apolipoprotein L family. Expressed in a variety of tissues, including liver, heart, uterus, spleen, colon, spinal cord, placenta, prostate and mammary gland, apoL-VI is thought to affect the movement of lipids in the cytoplasm and may allow the binding of lipids to organelles. Overproduction of apoL-VI induces apoptosis, suggesting that apoL-VI may also be involved in tumor progression. Like other members of the apolipoprotein L family, apoL-VI is thought to be involved in the development of schizophrenia.

REFERENCES

1. Dunham, I., et al. 1999. The DNA sequence of human chromosome 22. *Nature* 402: 489-495.
2. Page, N.M., et al. 2001. The human apolipoprotein L gene cluster: identification, classification, and sites of distribution. *Genomics* 74: 71-78.
3. Monajemi, H., et al. 2002. The apolipoprotein L gene cluster has emerged recently in evolution and is expressed in human vascular tissue. *Genomics* 79: 539-546.
4. Mimmack, M.L., et al. 2002. Gene expression analysis in schizophrenia: reproducible upregulation of several members of the apolipoprotein L family located in a high-susceptibility locus for schizophrenia on chromosome 22. *Proc. Natl. Acad. Sci. USA* 99: 4680-4685.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607256. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Liu, Z., et al. 2005. Apolipoprotein I6, a novel proapoptotic Bcl-2 homology 3-only protein, induces mitochondria-mediated apoptosis in cancer cells. *Mol. Cancer Res.* 3: 21-31.
7. Liu, Y.L., et al. 2008. RASD2, MYH9, and CACNG2 genes at chromosome 22q12 associated with the subgroup of schizophrenia with non-deficit in sustained attention and executive function. *Biol. Psychiatry* 64: 789-796.

CHROMOSOMAL LOCATION

Genetic locus: APOL6 (human) mapping to 22q12.3.

PRODUCT

apoL-VI (h2): 293T Lysate represents a lysate of human apoL-VI transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

apoL-VI (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive apoL-VI antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.