



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NALP1 (h2): 293T Lysate: sc-116236

BACKGROUND

NACHT-, LRR- and PYD-containing protein 1 (NALP1), also designated caspase recruitment domain protein 7, is a cytoplasmic protein. NALP1 contains a putative nucleotide binding site, a region of leucine-rich repeats, and death domain folds at both termini, providing protein/protein association functions such as caspase recruitment. NALP1 is involved in the innate immune response and is a component of the inflammasome. It forms cytoplasmic structures called death effector filaments and enhances APAF1 and cytochrome c-dependent activation of pro-caspase-9 and consecutive apoptosis. NALP1 is widely expressed in thymus, heart, spleen and peripheral blood leukocytes.

REFERENCES

- Hiller, S., Kohl, A., Fiorito, F., Herrmann, T., Wider, G., Tschopp, J., Grutter, M.G. and Wuthrich, K. 2003. NMR structure of the apoptosis- and inflammation-related NALP1 Pyrin domain. *Structure* 11: 1199-1205.
- Chamaillard, M., Girardin, S.E., Viala, J. and Philpott, D.J. 2003. Nods, NALPs and NAIP: intracellular regulators of bacterial-induced inflammation. *Cell. Microbiol.* 5: 581-592.
- Damiano, J.S., Oliveira, V., Welsh, K. and Reed, J.C. 2004. Heterotypic interactions among NACHT domains: implications for regulation of innate immune responses. *Biochem. J.* 381: 213-219.
- Sanz, C., Calasanz, M.J., Andreu, E., Richard, C., Prosper, F. and Fernandez-Luna, J.L. 2004. NALP1 is a transcriptional target for cAMP-response-element-binding protein (CREB) in myeloid leukaemia cells. *Biochem. J.* 384: 281-286.
- Liu, F., Lo, C.F., Ning, X., Kajkowski, E.M., Jin, M., Chiriac, C., Gonzales, C., Naureckiene, S., Lock, Y.W., Pong, K., Zaleska, M.M., Jacobsen, J.S., Silverman, S. and Ozenberger, B.A. 2004. Expression of NALP1 in cerebellar granule neurons stimulates apoptosis. *Cell. Signal.* 16: 1013-1021.

CHROMOSOMAL LOCATION

Genetic locus: NALP1 (human) mapping to 17p13.2.

PRODUCT

NALP1 (h2): 293T Lysate represents a lysate of human NALP1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

NALP1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive NALP1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

NALP1 (Nalpy1-4): sc-58550 is recommended as a positive control antibody for Western Blot analysis of enhanced human NALP1 expression in NALP1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



NALP1 (Nalpy1-4): sc-58550. Western blot analysis of NALP1 expression in non-transfected: sc-117752 (A) and human NALP1 transfected: sc-116236 (B) 293T whole cell lysates.



NALP1 (6D598): sc-71641. Western blot analysis of NALP1 expression in non-transfected: sc-117752 (A) and human NALP1 transfected: sc-116236 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.