



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CD100 (h): 293T Lysate: sc-116252

BACKGROUND

Semaphorins comprise a family of phylogenetically conserved proteins containing a 500 amino acid region termed the "Sema domain" in the amino-terminus. The semaphorin family is subdivided into eight groups. CD100 (SEMA4D) is a transmembrane protein belonging to group IV of the semaphorin family. CD100 is expressed in both embryonic and adult kidney, heart, lung and neural tissues. Plexin-B1, a receptor expressed abundantly in fetal brain and kidney, binds CD100 with high affinity. CD100 is the only member of the semaphorin family that is expressed in the immune system. CD100 activates T cell proliferation and promotes B cell aggregation. CD100 regulates B cell response by negating an inhibitory signal from CD72 that normally diminishes B cell response. This signal is turned off when CD100 binds to CD72 and induces tyrosine dephosphorylation of CD72, followed by the dissociation of SHP-1 from CD72. The gene encoding human CD100 maps to chromosome 9q22.2.

REFERENCES

1. Bougeret, C., Mansur, I.G., Dastot, H., Schmid, M., Mahouy, G., Bensussan, A. and Bomsell, L. 1992. Increased surface expression of a newly identified 150 kDa dimer early after human T lymphocyte activation. *J. Immunol.* 148: 318-323.
2. Herold, C., Bismuth, G., Bensussan, A. and Bomsell, L. 1995. Activation signals are delivered through two distinct epitopes of CD100, a unique 150 kDa human lymphocyte surface structure previously defined by BB18 mAb. *Int. Immunol.* 7: 1-8.
3. Furuyama, T., Inagaki, S., Kosugi, A., Noda, S., Saitoh, S., Ogata, M., Iwahashi, Y., Miyazaki, N., Hamaoka, T. and Tohyama, M. 1996. Identification of a novel transmembrane semaphorin expressed on lymphocytes. *J. Biol. Chem.* 271: 33376-33381.
4. Hall, K.T., Bomsell, L., Schultze, J.L., Boussiotis, V.A., Dorfman, D.M., Cardoso, A.A., Bensussan, A., Nadler, L.M. and Freeman, G.J. 1996. Human CD100, a novel leukocyte semaphorin that promotes B cell aggregation and differentiation. *Proc. Natl. Acad. Sci. USA* 93: 11780-11785.
5. Yu, H.H. and Kolodkin, A.L. 1999. Semaphorin signaling: a little less per-Plexin. *Neuron* 22: 11-14.
6. Tamagnone, L., Artigiani, S., Chen, H., He, Z., Ming, G.I., Song, H., Chedotal, A., Winberg, M.L., Goodman, C.S., Poo, M., Tessier-Lavigne, M. and Comoglio, P.M. 1999. Plexins are a large family of receptors for transmembrane, secreted, and GPI-anchored semaphorins in vertebrates. *Cell* 99: 71-80.

CHROMOSOMAL LOCATION

Genetic locus: SEMA4D (human) mapping to 9q22.2.

PRODUCT

CD100 (h): 293T Lysate represents a lysate of human CD100 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CD100 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD100 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

CD100 (C-3): sc-390675 is recommended as a positive control antibody for Western Blot analysis of enhanced human CD100 expression in CD100 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CD100 (C-3): sc-390675. Western blot analysis of CD100 expression in non-transfected: sc-117752 (A) and human CD100 transfected: sc-116252 (B) 293T whole cell lysates.



CD100 (D-5): sc-166367. Western blot analysis of CD100 expression in non-transfected: sc-117752 (A) and human CD100 transfected: sc-116252 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.