



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



MICAL1 (h): 293T Lysate: sc-116300

BACKGROUND

MICAL1 (microtubule associated monooxygenase, calponin and LIM domain containing 1), also known as MICAL or NICAL, is a 1,067 amino acid protein that localizes to both the cytoplasm and the cytoskeleton and contains one LIM zinc-binding domain and one calponin-homology domain. Expressed in kidney, thymus, spleen, lung and testes, MICAL1 interacts with the SH3 domain of Cas-L and, via this interaction, is thought to function as a cytoskeletal regulator that connects Cas-L to intermediate filaments. MICAL1 also interacts with Rab 1B, plexin-A3 and Vimentin, further regulating cytoskeletal events and possibly playing a role in axonal repulsion. Three isoforms of MICAL1 exist due to alternative splicing events.

REFERENCES

1. Terman, J.R., Mao, T., Pasterkamp, R.J., Yu, H.H. and Kolodkin, A.L. 2002. MICALs, a family of conserved flavoprotein oxidoreductases, function in plexin-mediated axonal repulsion. *Cell* 109: 887-900.
2. Suzuki, T., Nakamoto, T., Ogawa, S., Seo, S., Matsumura, T., Tachibana, K., Morimoto, C. and Hirai, H. 2002. MICAL, a novel CasL interacting molecule, associates with Vimentin. *J. Biol. Chem.* 277: 14933-14941.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607129. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Weide, T., Teuber, J., Bayer, M. and Barnekow, A. 2003. MICAL1 isoforms, novel rab1 interacting proteins. *Biochem. Biophys. Res. Commun.* 306: 79-86.
5. Bayer, M., Fischer, J., Kremerskothen, J., Ossendorf, E., Matanis, T., Konczal, M., Weide, T. and Barnekow, A. 2005. Identification and characterization of Iporin as a novel interaction partner for rab1. *BMC Cell Biol.* 6: 15.
6. Fischer, J., Weide, T. and Barnekow, A. 2005. The MICAL proteins and rab1: a possible link to the cytoskeleton? *Biochem. Biophys. Res. Commun.* 328: 415-423.
7. Jin, X., Zhang, J., Dai, H., Sun, H., Wang, D., Wu, J. and Shi, Y. 2007. Investigation of the four cooperative unfolding units existing in the MICAL1 CH domain. *Biophys. Chem.* 129: 269-278.
8. Schmidt, E.F., Shim, S.O. and Strittmatter, S.M. 2008. Release of MICAL autoinhibition by semaphorin-plexin signaling promotes interaction with collapsin response mediator protein. *J. Neurosci.* 28: 2287-2297.

CHROMOSOMAL LOCATION

Genetic locus: MICAL1 (human) mapping to 6q21.

PRODUCT

MICAL1 (h): 293T Lysate represents a lysate of human MICAL1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

MICAL1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive MICAL1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.