



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



ARIH1 (h): 293T Lysate: sc-116302



BACKGROUND

ARIH1 (ariadne homolog), also known as ubiquitin conjugating enzyme E2 binding protein 1, ARI, HARI, HHARI (human homolog of *Drosophila* ariadne), MOP-6 (monocyte protein 6) or UBCH7BP (UBCH7 binding protein), is a 557 amino acid cytoplasmic protein. Expressed in a wide variety of tissues, ARIH1 contains two RING-type zinc fingers and one IBR (in-between RING fingers)-type domain. ARIH1 is believed to be involved in protein degradation and protein translation. ARIH1 interacts with UBCH7 and is thought to function as an E3 ubiquitin-protein ligase (or as a component of an E3 complex) that, characteristic of E3 ligase proteins, accepts ubiquitin (in the form of a thioester) from an E2 ubiquitin-conjugating enzyme (UBCH7) and transfers that ubiquitin residue to substrates targeted for degradation. Specifically, ARIH1 interacts with and polyubiquitylates eIF4E2, thereby targeting it for proteasomal degradation.

REFERENCES

1. Moynihan, T.P., et al. 1999. The ubiquitin-conjugating enzymes UBCH7 and UBCH8 interact with RING finger/IBR motif-containing domains of HHARI and H7-AP1. *J. Biol. Chem.* 274: 30963-30968.
2. Tan, N.G., et al. 2000. Characterisation of the human and mouse orthologues of the *Drosophila* ariadne gene. *Cytogenet. Cell Genet.* 90: 242-245.
3. Ardley, H.C., et al. 2001. Features of the Parkin/ariadne-like ubiquitin ligase, HHARI, that regulate its interaction with the ubiquitin-conjugating enzyme, UBCH7. *J. Biol. Chem.* 276: 19640-19647.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605624. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Tan, N.G., et al. 2003. Human homologue of ariadne promotes the ubiquitylation of translation initiation factor 4E homologous protein, 4EHP. *FEBS Lett.* 554: 501-504.
6. Capili, A.D., et al. 2004. Structure of the C-terminal RING finger from a RING-IBR-RING/TRIAD motif reveals a novel zinc-binding domain distinct from a RING. *J. Mol. Biol.* 340: 1117-1129.
7. Okui, M., et al. 2005. Transcription factor single-minded 2 (SIM2) is ubiquitinated by the RING-IBR-RING-type E3 ubiquitin ligases. *Exp. Cell Res.* 309: 220-228.
8. Wong, E.S., et al. 2007. Relative sensitivity of Parkin and other cysteine-containing enzymes to stress-induced solubility alterations. *J. Biol. Chem.* 282: 12310-12318.
9. Schlehe, J.S., et al. 2008. Aberrant folding of pathogenic Parkin mutants: aggregation versus degradation. *J. Biol. Chem.* 283: 13771-13779.

CHROMOSOMAL LOCATION

Genetic locus: ARIH1 (human) mapping to 15q24.1.

PRODUCT

ARIH1 (h): 293T Lysate represents a lysate of human ARIH1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

ARIH1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive ARIH1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.